

- Jackson CL, de Jong IC. Clinicians and guidelines: leading a horse to water. *Med J Aust* 2001; 174: 267-268.
- Cameron B. The impact of pharmacy discharge planning on continuity of care. *Can J Hosp Pharm* 1994; 47: 101-109. □

### Catherine L Wilson

Project Officer, Southern Health, Monash Medical Centre, 246 Clayton Road, Clayton, VIC 3168  
catherine.wilson@med.monash.edu.au

**TO THE EDITOR:** It is refreshing to see quality initiatives like that of Mant et al,<sup>1</sup> which examine issues of continuity across different territories. Use of a simple audit tool (minimum dataset) and methodology has worked well to illuminate what misleadingly appears to be a simple problem (ie, the two-way exchange of information between the hospital and general practitioner in relation to medication). Unfortunately, like many problems that appear straightforward and easily fixable, the reality is that this issue is far more complex to address. A lack of clinical governance has been made apparent in both the public hospitals and the Divisions of General Practice that participated. Clinical governance demands that organisations be accountable for standards and performance in relation to clinical care,<sup>2</sup> and this is integral to addressing problems both internally and across the continuum of care.

Mant et al demonstrated that many hospitals had policies and strategies to accomplish the exchange of medication information,<sup>1</sup> indicating that these procedures were thought to be achievable with current resources. Before this study, knowledge among staff of implementation was scant and confused, and there was no system of review to reveal existing problems. When problems were exposed and changes made, staff were not upskilled to incorporate the change into their practice (eg, junior doctors not completing the new discharge referral form). Similarly, the Divisions of General Practice did not resolve issues surrounding the production and distribution of business cards that they had agreed to undertake.

This study has determined a means to measure performance and has intervened to analyse problems and yield some improvements. However, if the organisations involved do not commit to develop clinical governance, the path towards continued improvement will be extremely slow (if at all) and the findings of this project made irrelevant.

- Mant A, Kehoe L, Cockayne N, et al. A Quality Use of Medicines program for continuity of care in therapeutics from hospital to community. *Med J Aust* 2002; 177: 32-34.
- George M, Davey P. The responsibility of health boards for clinical governance. *The Quarterly* (Journal of the Royal Australasian College of Medical Administrators) 1999; 33(2): 20. □

### Peter W New

Head, Spinal Rehabilitation Unit, Caulfield General Medical Centre, 260 Kooyong Road, Caulfield, VIC 3162 p.new@cgmc.org.au

**TO THE EDITOR:** Mant et al<sup>1</sup> explore an important aspect of the quality use of medicines in their study on the continuity of medicines from hospital to community. Their study on compliance with an agreed minimum dataset for patient medication information exchange between hospitals and general practitioners provides a useful perspective of an approach to systems change. I wish to point out a number of limitations that may have affected their results and make some suggestions to improve the quality use of medicines.

GPs were audited on whether they provided medication information to hospitals. Many GPs work part-time. There is the possibility that the medical practice was contacted by a hospital employee, who obtained the information from a doctor other than the patient's usual GP.

The audit covered discharge summaries received by the GP by fax. Although faxing discharge summaries is convenient, there are potential problems with this method. There are the possibilities of dialling a wrong number, and faxed discharge summaries (particularly handwritten ones) may be difficult to read, which could also result in medication errors. In addition, a discharge summary may have been posted to the practice instead of faxed, which would under-report the true percentage of GPs who received the information.

It is not uncommon for patients to have multiple GPs.<sup>2</sup> However, it is my experience that only one GP is documented in the patient's medical file. This issue could have influenced the results of the GP audit and would be a further factor complicating the continuity of medicines from hospital to the community.

The authors mention the introduction of GP liaison officers to facilitate the notification of GPs about patient admissions and the rationale for medication changes. They do not report any other measures that they plan to introduce to improve their results. Given that systems problems have multifaceted answers, further expansion on what other steps could be taken would have been a useful addition to their article. I suggest that it would have been appropriate to include a broader range of key stakeholders in the workshops, such as community pharmacists and patients. In addition, a computerised hospital prescribing system could be integrated with an on-line evidence-based clinical guide to prescribing to assist in optimal medication selection. This could also be used to generate a discharge

medication list that was automatically sent to the patients' GPs. Such an approach would reduce errors and improve outcomes.<sup>3,4</sup>

- Mant A, Kehoe I, Cockayne NL, et al. A Quality Use of Medicines program for continuity of care in therapeutics from hospital to community. *Med J Aust* 2002; 177: 32-34.
- Veale BM, McCallum J, Saltman DC, et al. Consumer use of multiple general practitioners: an Australian epidemiological study. *Fam Pract* 1995; 12: 303-308.
- Bates DW, Leape LL, Cullen DJ, et al. Effects of computerized physician order entry and a team intervention on prevention of serious medical error. *JAMA* 1998; 280: 1311-1316.
- Monane M, Matthias DM, Nagle BA, Kelly MA. Improving prescribing patterns for the elderly through an online drug utilization review intervention: a system linking the physician, pharmacist, and computer. *JAMA* 1998; 280: 1249-1252. □

### Andrea Mant,\* Karen I Kaye,† Linda Kehoe,‡ Wendy C Rotem§

\*Adviser, Quality Use of Medicines Services, South East Sydney Area Health Service, Level 5, 376 Victoria Street, Darlinghurst, NSW 2010; †Executive Officer, NSW Therapeutic Assessment Group Inc., Sydney, NSW; ‡Consultant, Total Research, Sydney, NSW; §Consultant, Health Services Project Management, Sydney, NSW a.mant@unsw.edu.au

**IN REPLY:** We were pleased at the number of letters received exploring issues raised by our recent article.<sup>1</sup>

Jefford and colleagues question whether staff are aware of the Australian Pharmaceutical Advisory Council guidelines.<sup>2</sup> We expect to answer this question during the consultancy that the NSW Therapeutic Assessment Group is conducting for the Commonwealth Department of Health and Ageing. In this consultancy, we are evaluating the implementation and effectiveness of these guidelines. After its completion in October this year, we will have an implementation map of activities being undertaken relevant to the guidelines in Australia. Recommendations formalised at a national workshop will be made to the Department of Health and Ageing regarding future implementation, including alternative models and workable solutions.

All three letters raise questions about the key stakeholders for continuity of care. We invited a range of clinical and administrative staff, including senior and junior medical staff, nurse clinicians, clinical pharmacists and general practitioners from the relevant Divisions. However, participation of junior medical staff was limited. As Jefford et al comment, consumers and community pharmacists warrant further consideration in strategies for improving information exchange between hospitals and GPs. We also agree that patients' knowledge of their medications is often deficient: with limited resources, we were only able to address this through the GPs who interviewed their patients following discharge.