

practice improvement is needed. Improving the outcome of first stroke and reducing the incidence of second events with optimal secondary prevention would have clear benefits for the health of the nation.

*Stroke in Hospitals: an Australian Review of Treatment (START) investigators

START investigators other than the authors are:

Richard S Boyle (Head of Neurology, Princess Alexandra Hospital, Brisbane, QLD).

Richard P Gerraty (Neurologist, and Director, Stroke Unit, St Vincent's Hospital, Melbourne, VIC)

Graeme J Hankey (Head, Stroke Unit, Royal Perth Hospital, Perth, WA).

Peter Landau (Geriatrician, and Director, Stroke Service, Western Sydney Area Health Service, Sydney, NSW).

Stephen J Read (Neurologist, and Coordinator, Stroke Unit, Royal Brisbane Hospital, Brisbane, QLD).

John D G Watson (Neurologist, Royal Prince Alfred Hospital, Sydney, NSW).


COMPETING INTERESTS

Servier Laboratories supported this work financially but had no influence on the results or analyses presented.

Correction

Re: "Experimental human infection with the dog hookworm, *Ancylostoma caninum*", the Research article by Juergen K Landmann and Paul Prociv in the 20 January issue of the Journal (*Med J Aust* 2003; 178: 69-71), in which the labels for the second oral infection and cutaneous infection in the Figure in Box 2 were accidentally reversed. The corrected Figure is reproduced here.

2: Peripheral blood eosinophil levels after exposure to infective larvae of *Ancylostoma caninum*



* The three separate experiments were run consecutively over a period of one year.