

Improving access to acute stroke therapies: a controlled trial of organised pre-hospital and emergency care

Debbie A Quain, Mark W Parsons, Allan R Loudfoot, Neil J Spratt, Malcolm K Evans, Michelle L Russell, Angela T Royan, Andrea G Moore, Ferdinand Miteff, Carolyn J Hullick, John Attia, Patrick McElduff and Christopher R Levi

Care in a stroke unit improves long-term outcomes for stroke patients, underpinning the enhanced organisation of stroke services now underway across Australia.^{1,2} An important component of organised acute stroke care is thrombolytic therapy with intravenous tissue plasminogen activator (tPA),^{2,3} which was licensed for use in Australia in 2003. This highly cost-effective therapy⁴⁻⁶ substantially reduces major disability when used in an expert setting.^{7,8}

Only a small proportion of stroke patients currently receive thrombolytic therapy.⁹⁻¹¹ A major factor limiting patient access is that few arrive at expert stroke centres within the narrow 3-hour treatment window. Patient access to tPA could potentially be improved by extending organised stroke care through a cooperative system that crosses the traditional clinical and administrative boundaries of the acute stroke unit, emergency department (ED) and pre-hospital sector.

Where extended organised acute care has been successfully implemented, up to 20% of patients with ischaemic stroke are treated with tPA.¹²⁻¹⁴ Collaboration with ambulance services is recognised as a key to decreasing the time to presentation for patients with acute stroke. Pre-hospital stroke screening tools, such as the Los Angeles Prehospital Stroke Screen (LAPSS)¹⁵ and the Face Arm Speech Test (FAST),¹⁶ aid this process. These tools have acceptable diagnostic accuracy in the field and are important components of protocols for pre-hospital stroke care.

This study tested the effectiveness of an acute stroke care intervention developed in the Hunter Region of New South Wales — the Pre-hospital Acute Stroke Triage (PAST) protocol. This comprises a pre-hospital stroke assessment tool, an ambulance hospital-bypass protocol for thrombolysis-eligible patients outside the primary catchment area of the John Hunter Hospital, and the pre-notification and rapid deployment of the hospital's multidisciplinary stroke team. The PAST protocol was designed to reduce pre-hospital and ED delays to tPA treatment, and to streamline the initial management of patients by the existing acute stroke team.

ABSTRACT

Objective: To assess the effectiveness of the PAST (Pre-hospital Acute Stroke Triage) protocol in reducing pre-hospital and emergency department (ED) delays to patients receiving organised acute stroke care, thereby increasing access to thrombolytic therapy.

Design: Prospective cohort study using historical controls.

Setting: Hunter Region of New South Wales, September 2005 to March 2006 (pre-intervention) and September 2006 to March 2007 (post-intervention).

Participants: Consecutive patients presenting with acute stroke to a regional, tertiary referral hospital.

Intervention: PAST protocol, comprising a pre-hospital stroke assessment tool for ambulance officers, an ambulance protocol for hospital bypass for potentially thrombolysis-eligible patients, and pre-hospital notification of the acute stroke team.

Main outcome measures: Proportion of patients who received intravenous tissue plasminogen activator (tPA), process of care time points (symptom onset to ED arrival, ED arrival to tPA treatment, and ED transit time), and clinical outcomes of patients treated with tPA.

Results: The proportion of ischaemic stroke patients treated with tPA increased from 4.7% (pre-intervention) to 21.4% (post-intervention) ($P < 0.001$). Time point outcomes also improved, with a reduction in median times from symptom onset to ED arrival from 150 to 90.5 min ($P = 0.004$) and from ED arrival to stroke unit admission from 361 to 232.5 minutes ($P < 0.001$). Of those treated with tPA, 43% had minimal or no disability at 3 months.

Conclusions: Organised pre-hospital and ED acute stroke care increases patient access to tPA treatment, which is proven to reduce stroke-related disability.

MJA 2008; 189: 429-433

METHODS

Development of the PAST protocol

John Hunter Hospital, Newcastle, is the principal tertiary referral hospital for the NSW Hunter Region, serving a population of about one million people. The hospital's acute stroke team comprises an on-call stroke neurologist and an acute stroke nurse, as well as ED medical and nursing staff, and radiology staff.

Development of the protocol was coordinated by a steering committee comprising members of the ED, the acute stroke team and the NSW Ambulance Service. It involved collation and review of baseline stroke care performance data from the hospital's acute stroke database for the calendar years 2005 and 2006, with a specific focus on factors affecting tPA eligibility. Major system barriers to organised care and tPA therapy were identified as:

- lack of awareness among ambulance officers of the evidence and requirements for rapid access to organised acute stroke care;

- delivery of potentially thrombolysis-eligible patients to hospitals within the Hunter Region not equipped to provide a stroke thrombolysis service; and
- delays in acute stroke team notification on arrival of potentially thrombolysis-eligible stroke patients in the ED.

Components of the PAST protocol

To resolve these problems, the steering committee developed firstly a pre-hospital stroke assessment tool for ambulance officers, and secondly a pre-hospital notification system.

The pre-hospital stroke assessment tool was adapted from the United Kingdom FAST tool,¹⁶ to complement the National Stroke Foundation FAST community awareness campaign and existing assessment tools used by NSW ambulance officers. The local GAS-T tool has four elements:

- G = glucose in the "normal" range (4-17 mmol/L);
- A = arm drift, grip strength;
- S = speech disturbance; and
- T = time since symptom onset (Box 1).

1 Pre-hospital Acute Stroke Triage (PAST) protocol

HUNTER NEW ENGLAND
NSW HEALTH



Ambulance Service
of New South Wales

Pre hospital Acute Stroke Triage: Assessment Tool

TIME OF SYMPTOM ONSET MUST BE LESS THAN 2 HOURS

Patient Surname: _____ Case No: _____ Date _____
Time of patient assessment : _____

	YES	NO	UNSURE
Time of onset of symptoms: _____ (less than 2 hours)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
If the patient wakes with a deficit or cannot talk, then the time is taken from the last time the patient was seen without deficit.			

Glucose:

Is the patients BSL inside of the normal range 4mmol-17mmol YES NO UNSURE
Recorded BSL _____mmol/l @ _____hours.

Arm:

Lift the patient's arms both outstretched at 90° to trunk.
Ask the patient to hold them in that position for 5 seconds.

Does one arm drift down or fall rapidly	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Is handgrip weak on the same side?	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Is the loss of power noted on the	LEFT <input type="checkbox"/>		RIGHT <input type="checkbox"/>

Speech:

Attempt to have the patient say " You can't teach an old dog new tricks".
Ask a relative or friend if speech appears normal
Ascertain if speech is slurred or patient has difficulty finding words.

Is there discernable new speech impairment? YES NO UNSURE

CRITERIA FOR STROKE THROMBOLYSIS

- ✓ Must be YES to all of the above.
- ✓ The symptom onset time is definitely within 2 hours
- ✓ Symptoms not improving
- ✓ The patient is more than 18 years old
- ✓ The patient is normally ambulant Not previously wheel chair or bed bound
- ✓ The patient has no history of seizures/epilepsy

If the patient meets the criteria for thrombolysis, follow the Stroke Intervention Protocol.

It is vital that every attempt is made to have a relative attend the hospital with the patient unless this will cause a delay in transport.

An important difference from the UK tool is the replacement of the more difficult facial asymmetry test, "F", with a simple finger-prick glucose test, "G". Hyperglycaemia is a relative contraindication to tPA, being associated with poor outcomes and higher rates of haemorrhagic transformation,¹⁷ while hypoglycaemia is a known stroke mimic and may have led to reduced diagnostic accuracy in the field. "T" was added to the tool to emphasise the importance of the 2-hour time window. Although tPA is licensed for treatment of patients with ischaemic stroke within 3 hours of symptom onset, the third hour was allocated to in-hospital pretreatment procedures, such as patient assessment, imaging and gaining patient assent.

The cost of setting up the PAST protocol was minimal as the acute stroke team was already in place. Members of this team provided after-hours training for ambulance officers, and ambulance educators incorporated

lectures on stroke and the GAS-T tool into in-service sessions. Pocket cards outlining the GAS-T tool were funded by the NSW Ambulance Service Northern Division and distributed to all Hunter Region ambulance officers.

In addition, a pre-hospital notification system was set up, whereby the ambulance operations centre used Short Message Service (SMS) messaging to notify the on-call stroke neurologist when a patient met the GAS-T criteria. This allowed the neurologist to check the mandatory criteria before giving approval for the ambulance to bypass other regional hospitals when the patient was outside the primary catchment area of John Hunter Hospital. It also enabled rapid deployment of the acute stroke team to the ED to meet the ambulance at triage.

Evaluation of the PAST protocol

The study was a prospective cohort study using historical controls from the same

hospital in the corresponding period 1 year earlier. The study population comprised all patients with an initial diagnosis of acute stroke presenting to John Hunter Hospital. The intervention (PAST) group were those patients who presented to the hospital in the 6-month period from September 2006 to March 2007, immediately after implementation of the PAST protocol. The control group were those patients who presented in the corresponding period a year earlier, September 2005 to March 2006. Apart from the PAST protocol, the operating policies and procedures used by the acute stroke team were standard and did not alter over the period of the PAST protocol evaluation.

Outcome measures

The primary outcome measure was the proportion of ischaemic stroke patients who received intravenous tPA therapy in the PAST versus the control group.

Secondary outcome measures were process of care measures, including time from symptom onset to arrival at hospital (onset to door); time from arrival at hospital to commencement of tPA (door to needle); and transit time in the ED (door to admission to the ward, defined as the acute stroke unit). For the time-process measures, we compared patients with a definite symptom onset time within 24 hours of presentation to hospital in the PAST and control groups.

Tertiary outcome measures were based on clinical outcome at 3 months measured by the modified Rankin score (a validated disability measure where 0–1 = excellent outcome; 2 = functional independence; 3 = moderate disability; and 4–5 = major disability).^{18,19} Outcomes were compared between patients treated with tPA in the PAST group, and patients in the NINDS (National Institute of Neurological Disorders and Stroke) tPA trial⁷ and the SITS (Safe Implementation of Thrombolysis in Stroke) tPA registry.²⁰

Statistical analyses

Statistical analyses were performed using Stata version 9 (Stata Corp, College Station, Tex, USA). Continuous data were analysed using the Student *t* test when the assumption of normality appeared appropriate, and the Wilcoxon rank sum test when we believed this assumption was invalid. Categorical data were analysed using χ^2 tests or the Fisher exact test when the expected count in table cells was too low. *P* values less than 0.05 were considered statistically significant.

The Hunter New England Human Research Ethics Committee considered PAST as a quality improvement project, and individual patient consent was thus not required.

RESULTS

During the 6-month PAST protocol period, 232 patients presented to the John Hunter Hospital ED with an initial diagnosis of stroke, and 140 of these (60%) had a final

discharge diagnosis of ischaemic stroke (Box 2). Of the 232 patients, 122 (53%) had a defined onset time within the previous 24 hours, and 51 of the 232 were delivered to John Hunter Hospital using the PAST protocol.

In the 6-month control period, 205 patients presented to the John Hunter Hospital ED with an initial diagnosis of stroke, and 107 (52%) had a final discharge diagnosis of ischaemic stroke (Box 2). Of the 205 patients, 61 (30%) had a defined onset time within the previous 24 hours.

Access to tPA therapy

There was a statistically significant increase in the proportion of ischaemic stroke patients who received intravenous tPA therapy in the PAST group compared with the control group: 21.4% (95% CI, 14.6%–28.2%) versus 4.7% (95% CI, 1.5%–10.6%); $P < 0.001$ (Box 3). Expressed as a percentage of all patients presenting to the ED with an initial diagnosis of stroke, 12.9% (95% CI, 8.6%–17.2%) received tPA during the PAST period compared with 2.4% (95% CI, 0.8%–5.6%) during the control period ($P < 0.001$).

Among the 51 patients in the intervention period who met the PAST protocol criteria, 39 had a discharge diagnosis of ischaemic stroke, and 24 of the 39 were treated with tPA. A further six who did not meet the PAST protocol criteria for pre-notification received tPA, resulting in a total of 30 patients in the PAST period receiving intravenous tPA.

Among the 15 ischaemic stroke patients in the PAST group who did not receive tPA, stroke severity was considered too mild, or the deficits were rapidly resolving in six, another six had relative or absolute contraindications to tPA therapy, and the acute stroke team determined an earlier symptom onset time, outside the thrombolysis time window, for the remaining three.

In the control group, 61 patients (30%) arrived at the John Hunter Hospital ED within 24 hours of symptom onset, 38 of whom (62%) had a discharge diagnosis of ischaemic stroke. Five of these 38 received thrombolytic therapy.

Process of care indicators

Process of care measures are shown in Box 4. The median time from symptom onset to hospital arrival was reduced by 1 hour in the PAST group compared with the control group (90.5 v 150 min; $P = 0.004$). The median ED transit time was also reduced in

2 Patient characteristics, by treatment group

	Control group (n = 205)	PAST group (n = 232)
Male	104 (51%)	120 (52%)
Mean age in years (range)	69 (21–100)	71 (24–98)
Ischaemic stroke		
Total anterior circulation infarct	31 (15%)	53 (23%)
Partial anterior circulation infarct	37 (18%)	47 (20%)
Lacunar infarct	24 (12%)	23 (10%)
Posterior circulation infarct	15 (7%)	17 (7%)
Intracerebral haemorrhage	37 (18%)	29 (13%)
Transient ischaemic attack	45 (22%)	37 (16%)
Non-stroke	16 (8%)	26 (11%)
Onset time available	64 (31%)	122 (53%)
Onset within previous 24 h (% of patients with data available)	61 (95%)	122 (100%)
Presentation within 2 h of onset (% of patients with data available)	26 (41%)	68 (56%)

PAST = Pre-hospital Acute Stroke Triage.

3 Number of patients with ischaemic stroke and number who received tPA, by treatment group

	Control group (n = 205)	PAST group (n = 232)	P
No. with ischaemic stroke	107 (52.1%)	140 (60.3%)	0.12
No. who received tPA among			
All patients presenting with stroke	5 (2.4%)	30 (12.9%)	< 0.001
Ischaemic stroke patients	5 (4.7%)	30 (21.4%)	< 0.001

tPA = tissue plasminogen activator. PAST = Pre-hospital Acute Stroke Triage.

4 Process measures for patients with a defined symptom onset time within 24 hours of presentation to hospital and for patients who received tPA (values are median and IQR)

	Control group	PAST group	P*
Sample size			
All patients	61	122	
Recipients of tPA	5	30	
Onset to door time (min)			
All patients	150 (93–339)	90.5 (63–185)	0.004
Recipients of tPA	98 (85–101)	63.5 (52–89)	0.14
Door to ward time (min)			
All patients	361 (204–569)	232.5 (154–388)	< 0.001
Recipients of tPA	184 (120–204)	121 (77–179)	0.20
Door to needle time (min)			
Recipients of tPA	89 (85–160)	91.5 (70–100)	0.40

IQR = interquartile range. PAST = Pre-hospital Acute Stroke Triage. tPA = tissue plasminogen activator.

*Wilcoxon rank sum test.

