



Appendix 3

This appendix was part of the submitted manuscript and has been peer reviewed. It is posted as supplied by the authors.

Appendix to: Buckley NA, Whyte IM, Dawson AH, Isbister GK. A prospective cohort study of trends in self-poisoning, Newcastle, Australia, 1987–2012: plus ça change, plus c'est la même chose. *Med J Aust* 2015; 202: Online first 27 April 2015. doi: 10.5694/mja mja14.01116.

Table S1. Number (% of all drugs) of drugs usually obtained on prescription involved in self poisoning, and measures of morbidity (length of stay [median and interquartile range], intensive care unit admission, ventilation, in-hospital death). Agents over represented in intensive care unit admissions (>30%) shown in bold.

Drug	N (%)	LOS (h)	ICU (%)	Ventilated (%)	Deaths (%)
Antidepressants (all)	4477 (13.0%)	18 (11.9-32.9)	1001 (22.4%)	713 (15.9%)	6 (0.1%)
MAOIs	185 (0.5%)	18.4 (11.5-30.8)	31 (16.8%)	18 (9.7%)	0
SSRIs	1683 (4.9%)	15.3 (9.7-23)	158 (9.4%)	93 (5.5%)	0
SNRIs	780 (2.3%)	18.3 (12.3-29.6)	125 (16.0%)	74 (9.5%)	0
TCA s	1386 (4.0%)	24 (15.3-44.8)	635 (45.8%)	496 (35.8%)	6 (0.4%)
Antidepressants (Other)	443 (1.3%)	16.8 (11.6-23.6)	52 (11.7%)	32 (7.2%)	0
Antipsychotics	3180 (9.3%)	18.6 (12.5-33.5)	499 (15.7%)	331 (10.4%)	4 (0.1%)
Antipsychotics (Atypical)	1695 (4.9%)	18 (11.9-29.9)	259 (15.3%)	178 (10.5%)	2 (0.1%)
Antipsychotics (1st Generation)	1235 (3.6%)	18.5 (13-32.2)	198 (16.0%)	131 (10.6%)	2 (0.2%)
Lithium	250 (0.7%)	25.2 (14.9-69.6)	42 (16.8%)	22 (8.8%)	0
Anticonvulsants	1514 (4.4%)	19.4 (11.6-38)	253 (16.7%)	178 (11.8%)	2 (0.1%)
Barbiturates	28 (0.1%)	92.1 (39.3-140.9)	11 (39.3%)	15 (53.6%)	1 (3.6%)
Benzodiazepines	5470 (15.9%)	15.9 (9.8-25.3)	678 (12.4%)	449 (8.2%)	2 (0.04%)
Beta Blockers	289 (0.8%)	19.2 (12.6-36.3)	100 (34.6%)	39 (13.5%)	4 (1.4%)
Calcium Channel Blockers	139 (0.4%)	22 (14.4-43)	63 (45.3%)	9 (6.5%)	3 (2.2%)
Chloral Hydrate	23 (0.1%)	27.8 (20.2-41.9)	10 (43.5%)	8 (34.8%)	0
Colchicine	15 (0.04%)	69.8 (33.4-228)	9 (60.0%)	3 (20.0%)	2 (13.3%)
Dextropropoxyphene (± paracetamol/aspirin)	141 (0.4%)	15.8 (10.1-20.3)	17 (12.1%)	11 (7.8%)	0
Iron	39 (0.1%)	18.1 (13.9-27.1)	7 (17.9%)	5 (12.8%)	0
Theophylline	74 (0.2%)	28.5 (15.3-38.1)	40 (54.1%)	9 (12.2%)	1 (1.4%)
Sulphonylureas	34 (0.1%)	40 (21.7-71)	11 (32.4%)	3 (8.8%)	1 (2.9%)
Insulin	123 (0.4%)	37.2 (17.5-81.3)	42 (34.1%)	5 (4.1%)	2 (1.6%)
Other Prescribed Drugs ¹	3404 (9.9%)	-	-	-	-

¹ Only numbers of cases of all other prescribed drugs included.