



Appendix

**This appendix was part of the submitted manuscript and has been peer reviewed.
It is posted as supplied by the authors.**

Appendix to: Truswell S. Sceptics undermine effective dietary and heart health advice. *Med J Aust* 2015; 202: 412-414. doi: 10.5694/mja14.01448.

ON-LINE APPENDIX. References missed or incorrectly recorded in
CHOWDHURY et al *Annals of Internal Medicine* 2014; 160: 398-406

Prospective Studies

- [1] Laaksonen DE, Nyysonen K, Niksanen L, Rissanen TH, Salonen JT. Prediction of cardiovascular mortality in middle-aged men by dietary and serum linoleic and polyunsaturated fatty acids. *Arch Int Med* 2005; 165:193-199.
- [2] Ascherio A, Rimm EB, Giovannucci EL, Spiegelman D, Stampfer M, Willett WC. Dietary fat and risk of coronary heart disease in men: cohort follow up study in the United States. *Brit Med J* 1996; 313: 84-90.
- [3] Simpson HCR, Barker K, Carter RD, Cassels E, Mann JI. Low dietary intake of linoleic acid predisposes to myocardial infarction. *Brit Med J* 1982; 285: 683-684.
- [4] McGee DL, Reed DM, Yano K, Kagan A, Tillotson J. Ten-year incidence of coronary heart disease in the Honolulu Heart Program. *Am J Epidemiol* 1984; 119: 667-675.
- [5] Posner BM, Cobb JL, Belanger AJ, Cupples LA, D'Agostino RB, Stokes J. Dietary lipid predictors of coronary heart disease in men. The Framingham Study. *Arch Int Med* 1991; 151: 1181-1187.
- [6] Fraser GE, Sabate J, Beeson WL, Strahan TM. Protective effect of nut consumption on risk of coronary heart disease. The Adventist Health Study. *Arch Int Med* 1992; 152: 1416-1424.
- [7] Folsom AR, Arnett DK, Hutchinson RG, Liao F, Clegg LX, Cooper LS. Physical activity and incidence of heart disease in middle aged women and men. *Med Sci Sports Exerc* 1997; 29: 901-909.

Blood Lipid Fatty Acids

- [8] Pilz S, Scharnagl H, Tiran B, Wellnitz B, Seelhorst U, Boehm B, Marz W. Elevated plasma free fatty acids predict sudden cardiac death: a 6.85 year follow-up of 3315 patients after coronary arteriography. *Eur Heart J* 2007; 28: 2763-2769.
- [9] Lewis B. Composition of plasma cholesterol ester in relation to coronary heart disease and dietary fat. *Lancet* 1958; ii: 71-73.
- [10] Kirkeby K, Ingvaldsen P, Bjerkedal I. Fatty acid composition of serum lipids in men with myocardial infarction. *Acta Med Scand* 1972; 192: 513-519.
- [11] Salonen JT, Salonen R, Penttila I, et al. Serum fatty acids, apolipoproteins, selenium and vitamin antioxidants and the risk of death from coronary heart disease. *Am J Cardiol* 1985; 56: 226-231.

[12] Skuladottir G, Hardarson T, Sigfusson N et al. Arachidonic acid levels in serum phospholipids of patients with angina pectoris or fatal myocardial infarction. *Acta Med Scand* 1985; 218: 55-58.

[13] Valek J, Hammer J, Kohout M, Grafnetter D, Vondra K, Topinka V. Serum linoleic acid and cardiovascular death in postinfarction middle aged men. *Atherosclerosis* 1985; 54: 111-118.

[14] Block RC, Harris WS, Reid KJ, Spertus JA. Omega-6 and trans fatty acids in blood cell membranes: a risk factor for acute coronary syndromes. *Am Heart J* 2008; 156: 1117-1123.

Adipose Tissue Fatty Acids

[15] Kirkeby K, Nitter-Hauge S, Bjerkedal I. Fatty acid composition of adipose tissue in male Norwegians with myocardial infarction. *Acta Med Scand* 1972; 191: 321-324.

[16] Riemersma RA, Wood DA, Butler S et al. Linoleic acid content in adipose tissue and coronary heart disease. *Brit Med J* 1986; 292: 1423-1427.

[17] Wood DA, Riemersma RA, Butler S, Thomson M, Macintyre C, Elton RA, Oliver MF. Linoleic and eicosapentaenoic acids in adipose tissue and platelets and risk of coronary heart disease. *Lancet* 1987; i: 177-182.

[18] Roberts TL, Wood DA, Riemersma RA, Gallagher PJ, Lampe FC. Linoleic acid and risk of sudden cardiac death. *Brit Heart J* 1993; 70: 524-529.

[19] Pederson JI, Ringstad J, Almendingen K, Haugen TS, Steensvold I, Thelle DS. Adipose tissue fatty acids and risk of myocardial infarction—a case control study. *Eur J Clin Nutrition* 2000; 54: 618-625.

[20] Kark JD, Kaufman NA, Binka F, Goldberger N, Berry EM. Adipose tissue n-6 fatty acids and acute myocardial infarction in a population consuming a diet high in polyunsaturated fatty acids. *Am J Clin Nutrition* 2003; 77: 796-802.

[21] Clifton PM, Keogh JB, Noakes M. Trans fatty acids in adipose tissue and the food supply are associated with myocardial infarction. *J Nutr* 2004; 134: 874-879.

[22] Baylin A, Campos H. Arachidonic acid in adipose tissue is associated with nonfatal acute myocardial infarction in the Central Valley of Costa Rica. *J Nutr* 2004; 134: 3095-3099.