



## **Appendix**

**This appendix was part of the submitted manuscript and has been peer reviewed.  
It is posted as supplied by the authors.**

Appendix to: Möller H, Harvey L, Falster K, et al. Indigenous and non-Indigenous Australian children hospitalised for burn injuries: a population data linkage study. *Med J Aust* 2017; 206: 392-397. doi: 10.5694/mja16.00213.

**Appendix: Characteristics of burn injury and treatment by categories of geographical remoteness in a cohort of Aboriginal and non-Aboriginal children, NSW, 2000-2014.**

		Non-Aboriginal		Outer regional & Remote		Aboriginal		Outer regional & Remote	
		Major cities & Inner Regional	Percent	Number	Percent	Major cities & Inner Regional	Percent	Number	Percent
TBSA*	< 10% TBSA	3,116	87.2	430	85.7	138	84.7	112	79.4
	> 10% TBSA	458	12.8	72	14.3	25	15.3	29	20.6
Depth*	Superficial	249	7.1	48	9.6	17	10.8	10	8
	Partial thickness	2,737	78.3	368	73.9	116	73.4	88	70.4
	Full thickness	511	14.6	82	16.5	25	15.8	27	21.6
Severe burn <sup>#</sup>	No	3,282	88.7	477	87.7	156	89.1	121	81.8
	Yes	420	11.3	67	12.3	19	10.9	27	18.2
Severe burn treated at hospital with tertiary referral burn unit	No	178	42.4	38	56.7	7	36.8	12	44.4
	Yes	242	57.6	29	43.3	12	63.2	15	55.6
Anatomical location	Head/neck	693	18.7	113	20.8	39	22.3	29	19.6
	Trunk	800	21.6	96	17.6	28	16.0	38	25.7
	Shoulder/upper limb	493	13.3	76	14.0	19	10.9	16	10.8
	Wrist & hand	866	23.4	107	19.7	28	16.0	26	17.6
	Hip & lower limb	410	11.1	56	10.3	24	13.7	15	10.1
	Ankle & foot	286	7.7	61	11.2	24	13.7	13	8.8

\* Missing excluded

<sup>#</sup> Defined as partial or full thickness burn and > 10% TBSA