

Appendix 3

This appendix was part of the submitted manuscript and has been peer reviewed. It is posted as supplied by the authors.

Appendix to: Naylor JM, Hart A, Mittal R, et al. The value of inpatient rehabilitation after uncomplicated knee arthroplasty: a propensity score analysis. *Med J Aust* 2017; 207: 250-255. doi: 10.5694/mja16.01362.

Appendix 3

	Inpatient Group,	n =128*	No Inpatient Group, n = 129	
	%	No. of visits,	%	No. of visits,
		mean (sd)		mean (sd)
Receiving community	86.7	8.1 (5.2)#	74.0	5.9 (5.4)##
rehabilitation				
1-to-1 clinic	31.2	7.6 (3.9)	57.4	6.7 (4.8)
1-to-1 hydrotherapy	1.6	6.5 (0.7)	5.4	4.5 (3.3)
Group-based, clinic	9.3	8.0 (3.0)	7.0	5.3 (2.8)
Group-based,	4.7	3.0 (1.2)	3.1	5.8 (1.0)
hydrotherapy				
Domiciliary	1.6	2.0 (1.4)	10.1	7.8 (3.3)
Day program	53.1	9.7 (4.0)	4.7	12.8 (3.6)
*D (1 · · C 1	· · ·	•	•	

<i>Table A3. Profile of community-based physiotherapy rehabilitation</i>
--

*Details missing for 1 patient.

#Median 8 (IQR 5 to 11); ##Median 5 (IQR 0 to 9).