

Effectiveness of complementary and self-help treatments for anxiety disorders

Anthony F Jorm, Helen Christensen, Kathleen M Griffiths, Ruth A Parslow, Bryan Rodgers and Kelly A Blewitt

Anxiety disorders are among the most common mental disorders. They are estimated to affect 7% of men and 12% of women in Australia each year.¹ Anxiety disorders are also a major cause of disability: they account for around 2.7 million person-days out of role per month among Australian adults¹ and 6.5% of disability in the total population.²

In psychiatric classification systems, categorically distinct types of anxiety disorders are distinguished, such as panic disorder, obsessive-compulsive disorder, post-traumatic stress disorder and phobias. However, these anxiety disorders often occur together, and there is substantial comorbidity with depression and substance-use disorders.¹ While a categorical view of anxiety disorders may be useful for clinicians, community studies show a continuous dimension of anxiety severity, which is highly correlated with a dimension of depression severity.³ Disability increases as people move along this continuum.

There are effective medications and cognitive-behaviour treatments for anxiety disorders.⁴ However, only a fifth of people with anxiety as their principal complaint are estimated to have consulted a health professional in the previous 12 months.⁵ Furthermore, only half of these people received an intervention known to be effective for anxiety disorders.⁵ One reason people with anxiety disorders commonly give for not seeking professional help is that they prefer to manage the condition themselves.⁵ Indeed, people with a high level of anxiety and depression symptoms more commonly report using self-help methods than professional help.⁶

Australians also commonly use complementary therapies: an estimated 52% of Australian adults used complementary medicines in the year 2000, and 23% consulted complementary practitioners. These percentages have increased since 1993.⁷ Although we do not know how much of this use is for anxiety disorders, many of the complementary medicines available are marketed for "stress" or anxiety.

Given their frequent use, complementary and self-help treatments warrant the same level of evaluation as conventional treatments. The public needs to be informed about which treatments are effective, which are not, and which have not been adequately evaluated. General practitioners can play an important role in providing guidance. The purpose of our review is to provide GPs with an overview of the evidence on complementary

ABSTRACT

Objectives: To review the evidence for the effectiveness of complementary and self-help treatments for anxiety disorders.

Data sources: Systematic literature search using PubMed, PsycLit, and the Cochrane Library.

Data synthesis: 108 treatments were identified and grouped under the categories of medicines and homoeopathic remedies, physical treatments, lifestyle, and dietary changes. We give a description of the 34 treatments (for which evidence was found in the literature searched), the rationale behind the treatments, a review of studies on effectiveness, and the level of evidence for the effectiveness studies.

Conclusions: The treatments with the best evidence of effectiveness are kava (for generalised anxiety), exercise (for generalised anxiety), relaxation training (for generalised anxiety, panic disorder, dental phobia and test anxiety) and bibliotherapy (for specific phobias). There is more limited evidence to support the effectiveness of acupuncture, music, autogenic training and meditation for generalised anxiety; for inositol in the treatment of panic disorder and obsessive-compulsive disorder; and for alcohol avoidance by people with alcohol-use disorders to reduce a range of anxiety disorders.

MJA 2004; 181: S29–S46

and self-help treatments for anxiety disorders. We have previously reviewed the evidence on such treatments for depression.⁸

We distinguish between complementary treatment and self-help treatment:

- a complementary treatment is one involving practices and beliefs that are not generally upheld by the dominant health system in Western countries; and
- a self-help treatment is one that can be used by the public without necessarily consulting a health professional.
- While some self-help treatments are also complementary, others would be accepted by many practitioners in the dominant health system (eg, bibliotherapy, exercise). Each treatment is reviewed for its effectiveness in reducing generalised anxiety disorder or high levels of anxiety symptoms, as well as specific anxiety disorders (phobias, panic disorder/agoraphobia, obsessive-compulsive disorder, acute stress disorder, post-traumatic stress disorder).

METHODS

Our searches of the following sources identified 108 types of complementary and self-help treatments:

- clinical practice guidelines;^{4,9-19}
- brochures from health food shops;
- books and review articles on the topic;²⁰⁻²⁴ and
- key web sites — the top eight search engines, as rated by *searchenginewatch*, were used to compile a list of the top 20 Internet sites for each of the key search terms (given below),

Centre for Mental Health Research, Australian National University, Acton, ACT.

Anthony F Jorm, PhD, DSc, Director;

Helen Christensen, PhD, Deputy Director;

Kathleen M Griffiths, PhD, Director, Depression and Anxiety Consumer Research Unit;

Ruth A Parslow, PhD, Research Fellow;

Bryan Rodgers, PhD, Senior Fellow;

Kelly A Blewitt, BAppPsych, Research Assistant.

Correspondence: Professor Anthony F Jorm, Centre for Mental Health Research, Australian National University, Building 63, Eggleston Road, Acton, ACT 0200.

anthony.jorm@anu.edu.au

1 Modified* National Health and Medical Research Council (NHMRC) levels of evidence

Level	Description
I	Evidence obtained from a systematic review of all relevant randomised controlled trials.
II	Evidence obtained from at least one properly designed randomised controlled trial.
III-1	Evidence obtained from well-designed pseudorandomised controlled trials (alternate allocation or some other method).
III-2	Evidence obtained from comparative studies (including systematic reviews of such studies) with concurrent controls and allocation not randomised, cohort studies, case-control studies, or interrupted time series with a parallel control group.
III-3	Evidence obtained from comparative studies with historical control, two or more single-arm studies, or interrupted time series without a parallel control group.
IV	Evidence obtained from case-series, either post-test or pre-test/post-test.
V	No evidence, minimal evidence such as testimonials, single case studies, controlled studies of non-clinical samples only.

* The NHMRC evidence levels have been modified by adding a Level V to include even weaker types of evidence.

resulting in a list of about 110 sites, which were browsed to identify treatment recommendations.

Once the treatments were identified, PubMed, PsycLit and the Cochrane Library were searched using the following terms: name of treatment AND (anxiety OR panic OR phobic OR phobia OR agoraphobia OR post traumatic stress OR posttraumatic stress OR acute stress OR obsessive compulsive). The list of treatment terms used in the searches is available on request. The medical literature was searched up to 2 December 2002 for the earliest searches and 29 April 2003 for the latest ones. Articles were included if they reported treatment of individuals selected as having an anxiety disorder or a high level of anxiety symptoms. Articles on anxiety in normal samples were excluded. Occasionally, articles on anxiety symptoms in non-clinical samples (not selected for anxiety disorder or a high level of symptoms) are mentioned in the reviews below if they form an important part of the literature. However, they were not used in rating the effectiveness of treatments.

The evidence was evaluated using the National Health and Medical Research Council (NHMRC) levels of evidence modified by adding a Level V, as shown in Box 1.²⁵ We added a Level V to allow for even weaker types of evidence. It should be noted that these levels relate to the quality of the evidence, not the effectiveness of the intervention. A treatment could have been evaluated by rigorous methodologies and found to be ineffective or, conversely, evaluated by weaker methodologies, but found to be highly effective.

RESULTS

Treatments have been grouped under the following categories: medicines and homoeopathic remedies, physical treatments, lifestyle, and dietary changes. Most of the evidence relates to general-

ised anxiety disorder or high levels of anxiety symptoms, but when there is evidence relating to other anxiety disorders we flag this for each treatment. Information on safety is provided if available, but for many treatments there was none. For many treatments identified as being used for anxiety disorders, no evidence was found in the literature search. These treatments are listed in Box 2.

Medicines and homoeopathic remedies

Bach flower remedies

Description: Highly dilute tinctures of flowers are produced by adding particular flowers to spring water and preserving with alcohol.

Rationale: The Bach flower remedies are said to operate by transmitting energy from the flowers to the tincture.

Level of evidence: II.

Review of effectiveness: There have been two randomised double-blind, placebo-controlled studies of the effect of these remedies on test anxiety. In the first study, students who reported test anxiety were either administered the Bach “rescue remedy” or a placebo.²⁶ There was no consistent pattern to suggest that the rescue remedy reduced anxiety levels more than the placebo. The second study was a partial cross-over trial in which students were administered a mixture of 10 Bach flower remedies (impatiens, mimulus, gentian, chestnut bud, rock rose, larch, cherry plum, white chestnut, scleranthus and elm) or placebo.²⁷ There was no difference in anxiety reduction between the treatment and placebo groups.

Conclusion: Bach flower remedies do not appear to be an effective treatment for test anxiety. There is no evidence regarding the efficacy of these remedies for other anxiety disorders.

Berocca

Description: Berocca is a multivitamin preparation produced by Roche containing vitamin B complex, vitamin C, calcium and magnesium.

Rationale: This vitamin and mineral supplement may have stress-reducing effects.

Level of evidence: Level II.

Review of effectiveness: One randomised controlled trial (RCT) in healthy young adults²⁸ demonstrated reduced anxiety levels and improved psychological well-being in the intervention group relative to the group taking placebo, but there are no trials of individuals with high levels of anxiety or anxiety disorders.

Conclusion: There is no current evidence for an effect of Berocca on anxiety disorders.

Ginger (Zingiber officinale)

Description: A herbal remedy derived from a root and taken as tea or drink, or used as a tincture, in crystallised or tablet form.

Rationale: A traditional treatment for anxiety, it is now thought that ginger might reduce anxiety through antagonistic effects of its constituents (eg, gingerols) at the 5-HT₃ receptor.²⁹

Level of evidence: V.

Review of effectiveness: There are no studies of ginger administered alone for anxiety. Administration of *kampo* (ie, traditional Japanese) herbal preparations containing ginger was associated with reduced anxiety levels in four patients diagnosed as suffering from panic attacks with agoraphobia.³⁰ However, given the methodological limitations of case studies, and that the contribution of ginger

2 Treatments identified as being used for anxiety, but for which no evidence was found in the literature searched

Broad category	Treatments with no evidence
Medicines and homoeopathic remedies	American ginseng (<i>Panax quinquefolius</i>), ashwaganda (<i>Withania somnifera</i>), astragalus (<i>Astragalus membranaceus</i>), B vitamins, biotin, black cohosh (<i>Actaea racemosa</i> and <i>Cimicifuga racemosa</i>), borage (<i>Borago officinalis</i>), brahmi (<i>Bacopa monniera</i>), Californian poppy (<i>Eschscholtzia californica</i>), cat's claw (<i>Uncaria tomentosa</i>), catnip (<i>Nepeta cataria</i>), chamomile (<i>Anthemis nobilis</i>), chaste tree berry (<i>Vitex agnus castus</i>), Chinese medicinal mushrooms (reishi) (<i>Ganoderma lucidum</i>), choline, chromium, coenzyme Q10, cowslip (<i>Primula veris</i>), damiana (<i>Turnera aphrodisiaca</i>), dandelion (<i>Taraxacum officinale</i>), flax seeds (linseed) (<i>Linum usitatissimum</i>), GABA (γ -aminobutyric acid), ginkgo (<i>Gingko biloba</i>), ginseng (<i>Panax ginseng</i>), hawthorn (<i>Crataegus oxyacantha</i>), hops (<i>Humulus lupulus</i>), hyssop (<i>Hyssopus officinalis</i>), lecithin, lemon balm (<i>Melissa officinalis</i>), L-glutamine, L-tyrosine, melatonin, milk thistle (<i>Silybum marianum</i>), mistletoe (<i>Viscum album</i>), motherwort (<i>Leonurus cardiaca</i>), nettles (<i>Urtica dioica</i>), nicotinamide, oats (<i>Avena sativa</i>), omega 3 fatty acids, para-aminobenzoic acid (PABA), pantothenic acid, peppermint (<i>Mentha piperita</i>), phenylalanine, potassium, rehmannia (<i>Rehmannia glutinosa</i>), SAM-e, schizandra (<i>Schizandra chinensis</i>), selenium, Siberian ginseng (<i>Eleutherococcus senticosus</i>), skullcap (<i>Scutellaria lateriflora</i>), spirulina (<i>Spirulina maxima</i> and <i>S. platensis</i>), St Ignatius bean (<i>Ignatia amara</i>), taurine, tension tamer, tissue salts, vervain (<i>Verbena officinalis</i>), wild yam (<i>Dioscorea villosa</i>), wood betony (<i>Stachys officinalis</i> ; <i>Betonica officinalis</i>), yeast, zinc, zizyphus (<i>Zizyphus spinosa</i>).
Lifestyle	Adequate sleep, distraction techniques, marijuana (avoidance), pleasant activities, t'ai chi.
Dietary changes	Avoiding certain foods (barley, rye, wheat, dairy foods), sugar avoidance, ketogenic diet.

to the effects of the preparation is unknown, this study provides little useful information about ginger. It has been suggested that ginger might interact with warfarin to increase bleeding time,³¹ but a recent systematic review failed to confirm this potential interaction.³²

Conclusion: There is no scientifically reliable evidence as to the efficacy of ginger in the treatment of anxiety disorders.

Gotu kola (*Centella asiatica*)

Description: A traditional multipurpose herbal remedy used for anxiety and administered in the form of capsules, teas and tinctures.

Rationale: The constituents of gotu kola bind to cholecystokinin (CCK) receptors, which have been implicated in panic anxiety.³³ The herb may also affect anxiety through the GABA (γ -aminobutyric acid) system.³³ Gotu kola has been reported to decrease anxiety indicators (eg, acoustic startle response) in animals.³³

Level of evidence: V.

Review of effectiveness: There are no published studies of the effectiveness of gotu kola in the treatment of human anxiety disorders or symptoms of elevated anxiety. However, there is one study of the effect of gotu kola on anxiety symptoms in healthy participants. This small randomised double-blind, placebo-controlled study ($n=20$ per group) showed no change in self-rated anxiety half an hour to 2 hours after the administration of gotu kola, but acoustic startle response was attenuated at 60 and 90 minutes. The trial focused on the period immediately after a single administration of a large dose of gotu kola and did not use a standard measure of anxiety.

Conclusion: There is no evidence that gotu kola is useful for patients with anxiety disorders.

Combined preparations

Description: We defined combined preparations as mixtures containing a number of substances, including herbs. Combined preparations claiming to reduce anxiety include the Chinese medicine suanzaorentang, Worry Free (Maharishi Ayurvedic Prod-

ucts International, Colorado Springs, USA), Sedariston (a combination of St John's wort and valerian), and euphytose, a preparation commonly used in France.

Rationale: Claims of efficacy are based on traditional practice.

Level of evidence: II.

Review of effectiveness: Suanzaorentang has been tested in a time-series design in which a placebo was alternated with the herbal preparation in 30 individuals with reported high anxiety scores.³⁴ Anxiety scores decreased significantly after the week of active treatment and returned to previous high levels when taking the placebo. Worry Free has been tested in a pilot RCT in individuals with generalised anxiety disorder.³⁵ Ten patients were randomised to either Worry Free or placebo. After the 3-month trial period, those in the herbal preparation group had reduced symptoms and fewer had a diagnosis of anxiety disorder. Sedariston was compared with diazepam in a double-blind RCT of 100 outpatients with moderately severe anxiety disorders.³⁶ After 2 weeks, the group treated with Sedariston showed significantly larger decrements in trait anxiety scores than the group treated with diazepam. The effectiveness of euphytose has been tested on 182 outpatients with acute anxiety reactions, who were randomly assigned to euphytose or to placebo.³⁷ After 28 days, those taking euphytose had significantly reduced anxiety levels. It was well tolerated, and there were no differences in adverse events between the placebo and euphytose groups. However, it is thought that euphytose may potentiate an interaction effect with digoxin.³¹

Conclusion: There are too few studies to recommend the use of herbal preparations for treating anxiety. Suanzaorentang may be effective in anxiety, but it has not been evaluated in clinical groups or in RCTs. The pilot RCT of Worry Free produced good outcomes, but the number in each group was very small and the findings would need to be replicated. The one trial of Sedariston found that, compared with diazepam, it produced a greater reduction in anxiety. Euphytose may be effective in acute anxiety reactions, although its use in other anxiety disorders has not been evaluated.

Homoeopathy

Description: Homoeopathy uses preparations with highly dilute concentrations of a substance so that the molecules of the substance are not detectable.

Rationale: Homoeopathy assumes that symptoms reflect the body's attempt to heal itself. It is assumed that a substance that makes a well person ill with a particular set of symptoms will, in highly diluted form, promote recovery in an ill person who is suffering from the same set of symptoms.³⁸ Many theories have been proposed as to how homoeopathy might act on the body, including explanations based on quantum physics and information theory.³⁹ However, the precise mechanism is unknown.

Level of evidence: II.

Review of effectiveness: There have been two randomised double-blind, placebo-controlled studies of the effect of homoeopathy on anxiety. One study of primary care patients in whom either homoeopathy or a placebo was substituted for benzodiazepines reported no difference in anxiety between the homoeopathic-treated and placebo groups.⁴⁰ Another study of students with above-average anxiety levels reported less sleep loss but similar anxiety levels among a homoeopathic-treated group compared with a placebo group.⁴¹ However, given the multiple statistical comparisons involved, the positive result may have been achieved by chance.

Conclusion: There is no convincing evidence that homoeopathy is effective in treating anxiety.

Inositol

Description: Inositol is an isomer of glucose. Large amounts are present in the human body. It occurs in the normal human diet, with around 1 g/day being consumed.

Rationale: Inositol is involved in the phosphatidylinositol cycle, a second messenger system used by some serotonin and noradrenalin receptors.⁴²

Level of evidence: II.

Review of effectiveness: There have been five randomised double-blind trials of inositol at doses of 12–18 g/day. In panic disorder, inositol was found to be superior to placebo⁴³ and as effective as fluvoxamine.⁴⁴ In obsessive-compulsive disorder, it was found to be superior to placebo,⁴⁵ but did not augment the effects of selective serotonin reuptake inhibitors (SSRIs).⁴⁶ In post-traumatic stress disorder, however, inositol did not differ from placebo.⁴⁷ The size of these studies was small (13–21 patients) and treatment duration was short (4–6 weeks). In all studies, side effects were minimal.

Conclusion: There is some evidence for the effectiveness of inositol in treating panic disorder and obsessive-compulsive disorder. However, larger longer-term trials are needed.

Kava (Piper methysticum)

Description: The kava plant is a member of the pepper family. The root of the plant is used traditionally by Pacific Islanders to make a beverage that has social, ceremonial and medicinal uses. In Western countries, medicines containing kava extracts are marketed for the treatment of anxiety.

Rationale: The active ingredients, known as kava lactones, affect a range of neurotransmitter systems. Kava lactones are associated with the modulation of serotonergic and glutamatergic systems,

dopamine antagonism, and enhancement of the binding capacity of GABA receptors.⁴⁸

Level of evidence: I.

Review of effectiveness: A Cochrane review of 11 RCTs concluded that kava is superior to placebo for treating generalised anxiety.⁴⁹ Adverse events reported in these trials were mild, transient and infrequent. Kava is also non-addictive at therapeutic doses.⁵⁰ However, there have been reports of rare cases of liver failure in people taking high doses.⁵⁰ This has prompted authorities in some countries to take action — ranging from warnings to consumers to removing kava products from the market. The Australian Therapeutic Goods Administration has issued a voluntary recall of kava products, and consumers have been advised to discontinue use (for further information see <www.tga.health.gov.au/cm/kavafs0402.htm>).

Conclusion: Despite evidence of effectiveness of kava in treating general anxiety, rare cases of liver toxicity have led to advice against its use. Further data on its long-term safety are needed.

Lemongrass leaves (Cymbopogon citratus)

Description: Abafado is a herbal tea prepared from the dried leaves of lemongrass. It is used in Brazil.

Rationale: Lemongrass is believed to contain ingredients that reduce anxiety.

Level of evidence: II.

Review of effectiveness: One RCT of 18 individuals with high trait anxiety has been completed.⁵¹ Participants were provided with abafado or placebo, under double-blind conditions. Effects were observed 30 minutes later under mildly stressful conditions (the administration of a cognitive test). No differences in state anxiety scores were reported between the two groups.

Conclusion: Lemongrass in this form does not appear to have acute anxiolytic properties.

Licorice (Glycyrrhiza glabra)

Description: Licorice is extracted from the roots of this plant.

Rationale: Licorice contains compounds that inhibit monoamine oxidase.⁵²

Level of evidence: IV.

Review of effectiveness: The only evidence is from a small number of case studies claiming effectiveness.⁵³ Licorice should be used with caution, as excessive use can lead to hypertension, oedema and, in pregnancy, preterm birth.^{54,55}

Conclusion: There is no good quality evidence supporting licorice as effective in reducing anxiety.

Magnesium

Description: Magnesium supplements are available as tablets or an oral solution.

Rationale: It has been proposed that stress in people suffering from mental disorders can lead to marginal magnesium deficiency, which exacerbates symptoms such as anxiety.⁵⁶

Level of evidence: II.

Review of effectiveness: There have been no RCTs comparing magnesium with placebo in anxiety disorders. However, one RCT examined the effects of combining magnesium supplements with an anxiolytic medication over 10 days in women with mixed anxiety and depression.⁵⁷ Those receiving the anxiolytics plus

magnesium showed more rapid reduction in anxiety than a group receiving anxiolytics plus placebo.

Conclusion: There is little evidence at present on whether magnesium supplements are effective.

Passionflower (*Passiflora incarnata*)

Description: Passionflower is a plant native to North America. It is sold as a powder, a capsule, tablet or drops, and marketed as a treatment for sleep disorders, nervous tension and anxiety.

Rationale: Passionflower is a traditional remedy to treat anxiety. It is not known which constituents have sedative effects.⁵⁸

Level of evidence: II.

Review of effectiveness: A double-blind RCT in 32 outpatients with generalised anxiety disorder compared treatments of 45 drops of passionflower and 30 mg of oxazepam per day.⁵⁹ Four days after the trial began, there were no significant differences between the two treatments. Those taking passionflower reported fewer side effects than those taking oxazepam. There are no trials comparing passionflower with placebo.

Conclusion: While a small trial showed passionflower to be as effective as an anxiolytic, there is no evidence using placebo controls. From the evidence available it is not possible to say whether passionflower helps in the treatment of anxiety disorders.

St John's wort (*Hypericum perforatum*)

Description: St John's wort is a herb available as tablets, capsules or drops from supermarkets and health food shops.

Rationale: St John's wort is a traditional herbal remedy in Europe. Its mode of action is not fully understood, but it appears to inhibit the synaptic reuptake of serotonin, norepinephrine and dopamine.⁶⁰

Level of evidence: IV.

Review of effectiveness: While there is evidence supporting St John's wort as effective for depression,⁸ there are no RCTs in patients with anxiety disorders. The only evidence comes from an uncontrolled trial of patients with obsessive-compulsive disorder⁶¹ and case reports of patients with generalised anxiety disorder.⁶² The Therapeutic Goods Administration has warned that St John's wort may interact with a number of prescription medicines, leading to a loss of their therapeutic effect. Medicines affected include HIV protease inhibitors, HIV non-nucleoside reverse transcriptase inhibitors, cyclosporin, tacrolimus, warfarin, digoxin, theophylline, anticonvulsants, oral contraceptives, SSRIs and related drugs, and triptans. An information sheet for healthcare professionals is available at <www.tga.gov.au/docs/html/info.htm>.

Conclusion: There is only weak, uncontrolled evidence for the efficacy of St John's wort for treating anxiety disorders.

Valerian (*Valeriana officinalis*)

Description: The dried root of the valerian plant is a traditional herbal remedy, particularly in Hispanic cultures. Extracts are also available that contain some of the many ingredients. There are many species (probably over 200) of valerian, but the one most commonly used is *Valeriana officinalis*.

Rationale: Valerian is primarily used to treat sleep problems, but is also reported to have sedative properties. Many ingredients could contribute to this, including monoterpenes and sesquiterpenes, iridoids, alkaloids and amino acids.⁶³ Possible mechanisms include effects on the GABA system.

Level of evidence: II.

Review of effectiveness: One double-blind RCT allocated outpatients with generalised anxiety disorder to treatment with diazepam (2.5–7.5 mg/day), valepotriates (50–150 mg valerian extract per day) and placebo with 12 participants in each group.⁶⁴ Patients with comorbid anxiety and depressive disorders were excluded. All three groups showed significant decrements in rated and self-reported anxiety over the 4-week duration of the trial, but there were no significant differences between the three groups. The small number of participants and the relatively low doses of both diazepam and valerian extract were important limitations of the study. While the study had negative results, there is a report of valerian reducing psychic strain and feelings of somatic arousal in healthy subjects under stressful conditions, suggesting that further investigation is justified.⁶⁵ Valerian appears to have few side effects. There is one case report of an overdose,⁶⁶ and one report of a possible hazardous interaction with fluoxetine.⁶⁷ The reported hepatotoxicity may be due to other substances used in combination with valerian.⁶⁸ Some warnings have been given that valerian should not be used during pregnancy or lactation, and there have been more general recommendations not to use preparations based on Mexican (*Valeriana edulis*) or Indian (*Valeriana wallichii*) valerian.⁶⁹

Conclusion: There is little evidence regarding the treatment of anxiety disorders with valerian.

Vitamin C (ascorbic acid)

Description of treatment: Vitamin C is a water-soluble vitamin that is important in the formation of collagen and the absorption of iron.

Rationale: Ascorbic acid is said to modulate catecholaminergic activity and decrease stress reactions.

Level of evidence: V.

Review of effectiveness: The only evidence is from a case study in which vitamin C was used to treat obsessive-compulsive disorder.⁷⁰

Conclusion: There is insufficient evidence to evaluate the effects of ascorbic acid on anxiety symptoms.

5-hydroxy-L-tryptophan (5-HTP)

Description: This amino acid is naturally produced by the body and also available as a dietary supplement extracted from the plant *Griffonia simplicifolia*.

Rationale: A dysfunctional serotonergic system has been implicated in the development of some anxiety disorders. Since 5-HTP is the precursor of serotonin, treatment with 5-HTP could increase the level of serotonin production.

Level of evidence: II.

Review of effectiveness: One randomised, 8-week, double-blind, placebo-controlled trial compared the effect of 5-HTP (plus carbidopa), clomipramine and placebo in psychiatric outpatients with anxiety disorders without depression.^{71,72} The anxiety disorders included generalised anxiety disorder, agoraphobia and/or panic disorder, and obsessive-compulsive disorder. Compared with placebo, 5-HTP reduced anxiety levels. However, five participants dropped out due to increased anxiety, and results were only reported for those who completed the study.⁷² Another controlled study provided evidence that 5-HTP might prevent panic attacks. A double-blind placebo-controlled study of patients with panic disorder, challenged with CO₂ to induce anxiety, reported that

patients who had previously received a dose of 5-HTP developed fewer panic and anxiety symptoms than patients who had received placebo.⁷³ There have also been two small, open, uncontrolled studies of 5-HTP with long-standing anxiety disorders. In the first, nine of 10 patients with anxiety disorder (generalised anxiety disorder, or agoraphobia and/or panic disorder) showed improvement with the introduction of 5-HTP and carbidopa.⁷⁴ In the second study, five treatment-refractory patients with obsessive-compulsive disorder showed no reduction in obsessive-compulsive symptoms after an SSRI and pindolol were augmented with 5-HTP. The further addition of carbidopa to increase the uptake of 5-HTP in the brain created unacceptable side effects.⁷⁵ Elsewhere, it has been noted that 5-HTP should not be taken with SSRIs or monoamine oxidase inhibitors (MAOIs).⁷⁶ 5-HTP has also been found to contain contaminants associated with the eosinophilia-myalgia syndrome,⁷⁷ and the US Food and Drug Administration (2001) has raised concerns about the safety of “uncontaminated” 5-HTP in individuals who may be susceptible to eosinophilia-myalgia syndrome.⁷⁸

Conclusion: There is some evidence that 5-HTP might be effective in the treatment of anxiety disorders. However, until the risk of the eosinophilia-myalgia syndrome is clarified, 5-HTP cannot be recommended as a treatment for anxiety.

Physical treatments

Acupuncture

Description: Acupuncture is performed by stimulating designated points of the body through the insertion of needles, finger pressure, application of heat, or a combination of these treatments.

Rationale: According to traditional Chinese medicine, there is a network of energy (chi) that flows through the body in different channels. Any imbalance in this flow of energy creates a disease process, and acupuncture can correct such imbalances.

Level of evidence: II.

Conclusion: There is promising evidence that acupuncture can effectively reduce symptoms of anxiety in individuals with anxiety neuroses.

Review of effectiveness: One RCT has been conducted in which three groups, each comprising 80 individuals with confirmed anxiety neuroses, were treated with acupuncture only, behavioural desensitisation (including insight and supportive psychotherapy) only, or a combination of these treatments.⁷⁹ Those who received a combination of acupuncture and desensitisation showed significantly higher cure rates (52%) than those given either acupuncture only (20%) or desensitisation only (26%). For 60 of the 240 subjects followed up 1 year after treatment, the cure rates were 48%, 22% and 18%, respectively. Other studies reporting benefits of acupuncture have included only small numbers of patients receiving this treatment (fewer than 20).

Aromatherapy

Description: Aromatherapy involves exposure to the fragrances of essential oils from plants to promote healing. It is often delivered through skin massage or by means of a vaporiser.

Rationale: It is thought that aromas facilitate relaxation and exert a calming effect.

Level of evidence: II.

Review of effectiveness: There are no controlled studies of the effect of aromatherapy on patients with a clinical diagnosis of anxiety. One RCT of aromatherapy in men with speech anxiety found no difference in the anxiety reduction of men treated with aromatherapy and men in a control group.⁸⁰ A systematic review of RCTs of aromatherapy delivered through massage found that the intervention resulted in significantly lower anxiety levels than a control intervention in hospitalised cancer, cardiac and intensive care patients.⁸¹ However, participants were not specifically selected for anxiety, the reduction in anxiety was reported to be transient, and the trials contained methodological flaws.⁸¹

Conclusion: There is currently no evidence that aromatherapy is an effective treatment for anxiety disorders. However, it may provide short-term relief of anxiety symptoms in medical patients in hospital settings.

Hydrotherapy

Description: Hydrotherapy typically involves immersion in warm (close to body temperature) moving water.

Rationale: Hydrotherapy assists physical relaxation, but the exact mechanism for influencing anxiety is unknown.

Level of evidence: IV.

Review of effectiveness: One study of 14 patients with a range of anxiety disorders found a significant decrease in subjective anxiety and electromyograph readings from the frontalis muscles after 15 minutes of hydrotherapy with four water jets in a 38°C hot tub.⁸² No attempt was made to determine how lasting these changes were.

Conclusion: There is very little evidence for the effectiveness of hydrotherapy in treating anxiety disorders.

Massage/touch therapy

Description: Massage involves the smooth manual rubbing of muscle and other soft tissue, especially the back, shoulders and neck, usually by a trained masseuse or physiotherapist.

Rationale: Massage reduces cortisol, and possibly catecholamine, levels, although the precise mechanism behind this is not fully understood. Touch, itself, may be important, the therapy may promote a sense of being cared for, and physiological effects (eg, on vagal activity and on electroencephalogram recordings) may also be involved.^{83,84}

Level of evidence: II.

Review of effectiveness: One RCT of 60 children referred for signs of post-traumatic stress (in the aftermath of a hurricane) compared the effects of eight 30-minute sessions of back massage over a 1-month period with the effect of a similar number of control sessions of watching relaxing video movies.⁸⁵ Over the eight sessions, massage therapy was superior on several measures of anxiety. Other trials have examined effectiveness at reducing anxiety levels in other adolescent and adult patient groups (eg, with depression, bulimia and physical illnesses) and in non-patient samples.^{83,86-89}

Conclusion: There is some evidence that massage is effective in alleviating post-traumatic stress in children, but there is only very limited evidence for its effectiveness in adults or for treating other disorders. More studies of groups with anxiety disorders are warranted.

Lifestyle

Autogenic training

Description: Autogenic training is a self-relaxation procedure based on passive concentration on bodily perceptions (eg, heaviness and warmth of arms, legs and abdomen; rhythm of breathing and heartbeat).

Level of evidence: I.

Review of effectiveness: A meta-analysis of clinical outcome studies of autogenic training identified seven controlled studies that assessed the effectiveness of this treatment in reducing anxiety.⁹⁰ Three studies found that, compared with control groups, autogenic training significantly reduced anxiety for individuals with a diagnosed anxiety disorder, and those who described themselves as being anxious or having high levels of stress. One study found that student nurses given autogenic training for 6 weeks to reduce their risk of stress took significantly fewer days off compared with a control group. Two other studies found that autogenic training alone or autogenic training with other treatments had no significant effect on individuals' levels of test anxiety. The remaining study found that study participants who experienced "tension" reported that their levels of tension reduced by similar amounts after receiving no treatment, after receiving autogenic training only, or after receiving autogenic training and other treatments.

Conclusion: There is some evidence that autogenic training reduces levels of stress and anxiety symptoms. However, findings concerning autogenic training are mixed and further research needs to be undertaken.

Bibliotherapy

Description: Bibliotherapy is the use of written materials or computer programs, or listening to or viewing audiotapes or videotapes, for the purpose of gaining understanding or solving problems relevant to a person's developmental or therapeutic needs.

Level of evidence: I.

Review of effectiveness: A number of meta-analyses of the effectiveness of bibliotherapy in treating anxiety have been undertaken, the most recent being in 2003.⁹¹⁻⁹⁴ In these studies, bibliotherapy has been found to be most effective in reducing anxiety when the problem is circumscribed in its nature (eg, specific phobias), and when the individual is highly motivated to undertake treatment.⁹⁴ Bibliotherapy, by itself, had little effect on symptoms of panic disorder or obsessive-compulsive disorder.⁹⁴ Also, for most of these studies, the materials used were developed specifically for the research. The effectiveness of the many self-help materials that can be obtained by individuals in the community remains untested.

Conclusion: Bibliotherapy is most likely to reduce symptoms of anxiety associated with specific phobias. Further research is needed to determine which types of self-help tools are most effective and the conditions under which they are effective.

Dance and movement therapy

Description: Dance and movement therapies embrace a diverse range of treatments and activities, including professional help by trained therapists through to self-help forms of dance and movement. The therapy may be directly concerned with the individual's psychological problems or may take a form that has no overt link

to these problems. In some instances (eg, aerobic dance), dance is a medium for physical exercise (this is dealt with in the section on exercise (see page S36).

Rationale: Theories of movement therapy emphasise the importance of mind-body integration; the influence of physical movement on cognition, emotions and other behaviour; and the development of self-efficacy. Additional features include promotion of communication with others and reduction of feelings of isolation, although these are more relevant to treatment in residential settings.⁹⁵

Level of evidence: II.

Review of effectiveness: A small trial with severely disabled psychiatric outpatients (who had a variety of unreported diagnoses) compared changes in six patients who received movement therapy over 16 weeks with those in four control patients. Both groups also received medication and group therapy. Patients in the experimental group showed a significantly greater decrease in muscular tension and trait anxiety measures, but not in state anxiety. The report does not describe the method of allocating patients to the experimental and control groups.⁹⁶ A second trial, using randomised allocation, evaluated short-term changes in anxiety and depression after a single session of dance therapy for groups of hospitalised psychiatric patients ($n = 20$) and individuals drawn from the neighbourhood community ($n = 20$). The experimental groups (10 subjects in each) showed significantly greater decrements in anxiety and depression scores than the control group over the 1-hour period. The degree of change between the patient and community participants was not significantly different.⁹⁷ A further RCT of volunteer students with self-identified problems of test anxiety found that four 35-minute movement sessions over 2 consecutive weeks led to a significantly greater reduction in total scores for the Test Attitude Inventory in the experimental group ($n = 11$) compared with the control group ($n = 10$).⁹⁸ The sessions were described as "quasi-educational in nature, with a cognitive/behavioural and psychoeducational emphasis", and the content specifically addressed test anxiety. There are also five case reports of the effectiveness of Dosa therapy (a Japanese form of movement therapy) in reducing anxiety levels in female graduate students with anxiety problems.⁹⁹ Short- and long-term changes in anxiety associated with dance therapy and dance classes have also been studied in groups unselected for anxiety levels or disorders. One report found significantly greater reductions in questionnaire measures of anxiety in student psychiatric nurses who received a single session of dance therapy compared with those in the control group. This applied in one study in which 17 students were randomly allocated to dance therapy or to no intervention, and also in a second study in which 46 students were randomly allocated to dance therapy or verbal therapy.¹⁰⁰ An observational study of student classes in colleges of further education found a significant decrement of almost 1 SD in state anxiety scores over a term of study for 23 modern dance students.¹⁰¹ The change was greater than that found for students in sport, music and mathematics classes (all non-significant).

Conclusion: There is some evidence that structured movement therapy is helpful in reducing test anxiety, and it may also be helpful for managing other anxiety problems in clinical groups and in people with self-identified anxiety. The suggestion that conventional dance classes may alleviate state anxiety (more so than other physical exercise) is worthy of further study.

Exercise

Description: Exercise is physical activity, which may be aerobic (eg, running), with the aim of increasing cardiovascular fitness and stamina, or anaerobic (involving weight training), with the aim of increasing strength and endurance.

Rationale: Possible mechanisms include that exercise acts as a buffer for stress; that exercise engages the same processes as those elicited by meditation; that exercise might act to put people with agoraphobia into unsafe environments, and thereby offer “exposure therapy”; or that exercise might produce more monoamine neurotransmitters, which may reduce anxiety.¹⁰² Some authors have suggested that exercise is potentially more likely to be effective for some anxiety conditions, such as generalised anxiety disorder and panic disorder, but not for others, such as phobias and post-traumatic stress disorder (given that these conditions are associated with responses to specific situation cues or experiences).¹⁰³ It has also been suggested that there may be multiple and synergistic causes of exercise’s effectiveness, given that exercise affects biological and psychological processes.

Level of evidence: I.

Review of effectiveness: There have been numerous meta-analyses of the effect of exercise on anxiety symptoms. However, only two have examined the effects of exercise on individuals with elevated anxiety levels. In one meta-analysis of 104 studies, there were 11 that yielded data on trait anxiety for individuals who had been specifically identified as highly anxious.¹⁰⁴ The mean effect size from these comparisons was 0.47 SD units, indicating that, relative to results in comparison groups, exercise was associated with a moderate reduction in anxiety. The second meta-analysis examined the effect of aerobic and non-aerobic physical exercise on symptoms of anxiety and depression.¹⁰⁵ Individual controlled and randomised controlled trials were included if groups had anxiety scores above the 50th percentile. The latter meta-analysis identified 11 studies that compared an exercise group with a control group of patients waiting for treatment (wait-list controls) and revealed an effect size of 0.94. Within these comparisons, seven studies included individuals with a formal anxiety diagnosis, while the remainder consisted of individuals with higher than average anxiety. Studies of individuals with a diagnosis were associated with an effect size of 0.99, and those undiagnosed with an effect size of 0.85. The major limitation of this meta-analysis from our perspective is the inclusion of individuals with mild to moderate anxiety (50th percentile or above) who did not meet anxiety diagnostic criteria, and the lack of outcome data for different types of anxiety disorders. Moreover, of the studies with “a strictly defined group of participants with diagnoses of anxiety disorders”, there were none that compared the effect of exercise with a placebo control or with the effect of psychotherapy or medication.

However, outcomes for specific types of anxiety conditions have been reported for generalised anxiety disorder and high trait anxiety, with two controlled trials indicating aerobic exercise to be superior to strength and mobility exercises,¹⁰⁶ and a 6-week jogging intervention to be as effective as cognitive behaviour therapy and more effective than no treatment.¹⁰⁷ For panic disorder, Brooks et al compared a 10-week running program with either treatment with clomipramine or with a placebo.¹⁰⁸ At the end of the intervention, both active treatments were more effective than placebo, but clomipramine had an earlier effect, a stronger effect on three of the five outcome measures, and was associated with a lower drop-out rate (zero, compared with 31% for the

exercise group). Panic disorder has been reported to respond to both aerobic and non-aerobic interventions.¹⁰⁹ Finally, Manger examined the performance of 26 volunteers who met criteria for post-traumatic stress disorder.¹¹⁰ The intervention was 6–10 weeks of treadmill exercise for 30 minutes, two to three times a week. Only nine subjects remained after the intervention. Despite methodological problems, the authors suggested that the effects of exercise on post-traumatic stress disorder were promising.

Conclusion: Exercise has demonstrated strong effects for mild to moderate anxiety, but further studies are required to determine its effects on specific anxiety conditions. There is one trial that shows exercise to be as effective in the longer term as standard use of highly effective anti-anxiety medication (clomipramine) in panic disorder. Clearly, there is a need to investigate the potentially strong therapeutic effects of exercise on specific anxiety conditions, and further RCTs are required. Exercise is a highly promising treatment for anxiety.

Humour

Description: Humour allows us to see as amusing or ridiculous aspects of both everyday life and the unusual events that confront us.

Rationale: Laughter is claimed to be beneficial to the autonomic system. It can also provide a physical release for accumulated tension, and hence may moderate the effects of stressful life events on physical and psychological feelings.

Level of evidence: V.

Review of effectiveness: There have been a number of RCTs, all with undergraduate students, that have tested the effects of humour on symptoms of anxiety. Students treated for arachnophobia with either systematic desensitisation or humour-based desensitisation reported the same degree of reduction in their phobia. However, humour alone was not used in this study.¹¹¹ Others have examined the effects of humour on healthy students placed in anxiety-provoking situations. One such study found that having a sense of humour reduces the likelihood that experiences of stress will cause anxiety, but this occurred in men only.¹¹² In another trial, students who expected that they would soon experience an unpleasant event showed fewer signs of stress when they listened to a humorous tape while waiting compared with those who listened to a non-humorous tape or no tape.^{113,114} Similarly, those with a good sense of humour showed fewer symptoms of stress in such circumstances compared with those with a poor sense of humour.¹¹³

Conclusion: There is some indication that humour can reduce anxiety in undergraduate students in laboratory settings. However, there is no evidence on the use of humour in anxiety disorders.

Meditation

Description: Meditation involves focusing the mind on an object or word combined with a passive attitude and a comfortable physical position.

Rationale: Meditation is believed to produce a state of relaxation that is incompatible with feelings of anxiety to induce a “trance state” that elicits positive mood and/or to provide a period of “rest”.

Level of evidence: II.

Review of effectiveness: Five RCTs have examined the effect of meditation in generalised anxiety disorder or high trait anxiety.

Four of the five trials found that meditation produced equivalent effects to other forms of relaxation, including applied relaxation, progressive muscular relaxation and biofeedback,¹¹⁵⁻¹¹⁸ and these effects were superior compared with those in a control group (wait-list controls) but not superior to “sitting meditation”.^{116,118} By contrast, another study reported no difference between meditation, relaxation, an “anti-relaxation” control and no treatment control.¹¹⁹ Another trial compared meditation to relaxation, skills acquisition and practice-only for test anxiety and found that meditation was more effective than practice-only, but less effective on test performance than skills acquisition.¹²⁰ Shannahoff-Khalsa reported that a yogic meditation technique (based on kundalini yoga) was more effective than a standard relaxation procedure in the treatment of obsessive-compulsive disorder.¹²¹ There have also been case studies of the effects of meditation on dental fear.^{122,123}

Conclusion: Meditation may be an effective intervention for high trait anxiety and generalised anxiety disorder. Its effectiveness in treating other forms of anxiety disorder has not been established.

Music

Description: Music is widely used by the public for reducing anxiety, but also by music therapists, who may combine it with other therapeutic elements.

Rationale: Music is a traditional intervention.

Level of evidence: II.

Conclusion: There is some evidence to support the anxiety-reducing effects of music. However, music has only been evaluated in people with high levels of anxiety symptoms, not with diagnosed anxiety disorders.

Review of effectiveness: There have been three RCTs comparing music interventions with no treatment in people with high levels of anxiety symptoms. One study involved a single session and produced no improvement;¹²⁴ another involved three sessions and produced improvement at the end of treatment that was not maintained 6 months later;¹²⁵ the third trial involved 10 sessions and produced an improvement which was maintained 2 months later.¹²⁶ Music has also been shown to have the same degree of anxiety-reducing effect as muscle relaxation¹²⁷ and to boost the effect of a brief cognitive reframing intervention.¹²⁸

Prayer

Description: Prayer can be used to promote healing of oneself or of someone else (intercessory prayer). It can also be used as an adjunct to psychotherapy.

Rationale: Prayer has traditionally been used in time of illness and is often used by the public to help in coping with mental health problems.

Level of evidence: II.

Review of effectiveness: There have been no controlled trials of either personal or intercessory prayer in anxiety disorders. However, there has been one RCT adding a religious component, including prayer, to standard treatment for generalised anxiety disorder in religious patients.¹²⁹ This trial found that including a religious component produced greater improvement after 3 months' treatment, but no difference after 6 months.

Conclusion: There is currently no evidence to support prayer alone as a treatment for anxiety disorders.

Relaxation

Description: Relaxation therapy involves a family of techniques to elicit the relaxation response.¹³⁰ A variety of types are distinguished, including progressive relaxation, release-only relaxation, cue-controlled relaxation, differential relaxation, rapid relaxation and applied relaxation. Progressive relaxation, probably the most common form, teaches individuals to systematically identify and relax specific muscle groups.

Rationale: Relaxation induces a state that is incompatible with the physiological and psychological arousal associated with anxiety. Relaxation teaches individuals to recognise symptoms of anxiety and respond to them with a technique that reduces arousal.

Level of evidence: I or II.

Review of effectiveness

Generalised anxiety disorder and high trait anxiety: A meta-analysis reported relaxation to be superior by an average of 0.60 SD units relative to non-active interventions (for example, waitlist control).¹³¹ We identified 26 controlled trials^{116-118,127,132-153} in which relaxation was found to be generally superior to the control conditions, and to be as effective as a range of psychological treatments, including cognitive behaviour therapy, meditation, autogenic training and biofeedback. Anxiety management¹³⁶ and muscle stretching¹³⁹ were reported to be superior in one study each.

Panic disorder and agoraphobia: A meta-analysis identified relaxation to be as effective as non-active interventions, with an effect size of about 0.58.¹⁵⁴ Drug treatments were compared in two studies. These indicated an effect size of 0.22 in favour of the drug treatments. Five studies compared relaxation with other forms of behavioural therapy, including exposure therapy. These yielded an effect size of 0.08 in favour of behavioural therapies. These findings are consistent with our review of 11 controlled trials of relaxation therapy:¹⁵⁵⁻¹⁶⁵ in four, with a wait-list control group or placebo control, relaxation was better; relaxation was as effective as behavioural interventions, such as cognitive behaviour therapy or cognitive therapy (as effective in four of five studies, less effective in one), it was more effective than paradoxical intention (one study), as effective or almost as effective as exposure-based therapies in the short term (five studies, with one finding it was not as effective),¹⁵⁷ and roughly equivalent to drug treatments (two studies).

Post-traumatic stress disorder: A meta-analysis of treatments for post-traumatic stress disorder identified one study in which relaxation (effect size, 0.45) was inferior to behavioural therapies for this disorder (effect size, 1.27).¹⁶⁶ Our review of six studies¹⁶⁷⁻¹⁷² indicated that, although relaxation may be superior to no intervention (wait-list controls), it appears to be less effective than cognitive restructuring, social skills training, exposure therapy and combined treatments.

Obsessive-compulsive disorder: There have been four trials of relaxation therapies in obsessive-compulsive disorder.¹⁷³⁻¹⁷⁶ One involved students diagnosed with obsessive-compulsive disorder.¹⁷³ Relaxation was found to be inferior to behavioural therapy and in-vivo exposure therapy, respectively.^{174,175} Greist et al reported that relaxation was inferior to both computer-assisted and clinician-assisted behaviour therapy.¹⁷⁶

Social phobia: There have been two studies of social phobia:^{177,178} in one study with a wait-list control group, relaxation was reported to be superior; it was as effective as self-instructional training,¹⁷⁸

and less effective than social skills training.¹⁷⁷ These studies yielded a combined pre–post effect size of 0.51 (95% CI, 0.45–0.57), which was of smaller magnitude than effect sizes reported for benzodiazepines and SSRIs (effect sizes, 1.5) in a meta-analysis reported by Fedoroff and Taylor.¹⁷⁹

Dental phobia: Five controlled trials have identified that relaxation is as effective as other behavioural and drug interventions in dental phobia.^{180–184}

Small-animal phobia and specific phobias: Seven trials have indicated that, for snake or spider phobias, there is little evidence that relaxation is more effective than control, but that exposure-based therapies are effective.^{185–191} One study has indicated that both relaxation and stress inoculation treatment (a technique involving the reconstruction of negative thoughts while physically present in the feared situation) produce pre–post change in people with fear of flying. For claustrophobia, Ost et al reported that both applied relaxation and exposure produced change.¹⁹² Two studies of blood phobia indicated that applied relaxation provided a slight advantage which did not persist at follow-up.^{193,194}

Speech anxiety: Of eight RCTs, three (using wait-list controls or placebo) found relaxation to be more effective than, and three found it to be as effective as, other behavioural therapies such as biofeedback, systematic desensitisation, and speech practice.^{195–202} Rational emotive therapy has been found to be more effective,¹⁹⁷ but relaxation has been found to be superior to aromatherapy.²⁰⁰

Test anxiety: Of 29 RCTs, those using wait-list controls or other controls found that relaxation is more effective, and the rest found it to be as effective as other behavioural therapies for test anxiety, such as systematic desensitisation, rational emotive therapy, biofeedback, rehearsal and skills acquisition, and meditation.^{203–231}

Conclusion: Relaxation is an effective therapy for dental phobia, test anxiety, panic disorder and generalised anxiety disorder. In particular, there is no strong evidence to indicate that validated exposure-based or efficacious drug treatments are more effective than relaxation. However, although relaxation lowers symptoms relative to those in wait-list controls, it may be less effective than other behavioural treatments for small-animal and specific phobias, social phobia, post-traumatic stress disorder and obsessive-compulsive disorder.

Yoga

Description: Yoga includes exercises for attaining bodily and mental control and wellbeing.

Rationale: Yoga is often used for relief of stress and anxiety.

Level of evidence: II.

Review of effectiveness: One study showed that yoga was superior to diazepam for generalised anxiety, but patients were assigned to yoga treatment based on preference rather than random assignment.²³² In another randomised trial, test anxiety was treated with a set of yoga exercises combined with one of the following — autosuggestion, progressive muscle relaxation or a control talking session.²²⁷ Yoga treatment was found to be superior to both relaxation and control treatments on one outcome measure, but not on another. However, the results were not fully described and the outcome measures were of unknown validity.

Conclusion: In the absence of any well-conducted studies, it is impossible to say whether yoga is effective.

Dietary changes

Alcohol

Description: Alcohol is commonly used to reduce anxiety in the short term, but prolonged use may lead to greater anxiety. For this reason, it has been recommended that people with anxiety or other psychiatric disorders should avoid alcohol altogether.²³³

Rationale: Heavy drinkers have an increased risk of suffering from anxiety disorders. There are two main ways in which heavy drinking might lead to increased anxiety. Firstly, while acute alcohol intoxication affects GABA receptors in a similar way to benzodiazepines, chronic alcohol use leads to decreased GABA tone and generates anxiety.²³⁴ Secondly, heavy alcohol consumption leads to problems (eg, financial, occupational, relationship and health problems) which may increase anxiety.

Level of evidence: II (for the short-term anxiety-reducing properties of alcohol); IV (for the benefits of abstinence in heavy drinkers).

Review of effectiveness: There have been a number of controlled trials examining the short-term effects of alcohol on anxiety. In these trials, people with a range of anxiety disorders have been given alcohol or a placebo drink and then exposed to a stressful situation. Many of these studies show that alcohol reduces anxiety in response to the stressor,^{235–243} but there are negative studies as well.^{240–243} There have been no controlled trials of heavy drinkers with anxiety disorders abstaining from alcohol. However, uncontrolled studies of patients admitted to alcohol detoxification programs show very high levels of anxiety initially and a very rapid decline in anxiety symptoms after cessation of alcohol.^{244–249} The rapidity of the response suggests that alcohol cessation is producing this effect.

Conclusion: There is some evidence that use of alcohol has a short-term anxiety-reducing effect. However, for those with alcohol-use disorders, eliminating alcohol use may be an effective way of reducing anxiety.

Caffeine reduction

Description: Caffeine occurs naturally in the leaves, seeds and nuts of some plants. It is found in coffee, tea, soft drinks (eg, cola), chocolate, cocoa and some medications.

Rationale: It has been suggested that caffeine induces anxiety by binding to adenosine receptors and blocking the anxiolytic effects of adenosine.²⁵⁰

Level of evidence: IV (for caffeine reduction); II (for caffeine challenges).

Review of effectiveness: Several researchers have reported case studies or case series where caffeine reduction was associated with decreased anxiety levels among patients with anxiety disorders.^{251–254} In addition, in retrospective surveys, people with anxiety disorders have reported high rates of caffeine discontinuation due to perceived anxiety-producing side effects.^{255,256} However, there have been no controlled studies of the effect of caffeine abstinence or reduction on anxiety disorders. Clinical recommendations that patients with anxiety disorders should avoid caffeine are primarily based on the results of caffeine challenge studies. These studies measure the immediate effect on anxiety of the short-term administration of caffeine to people with anxiety disorders after a caffeine-free period. Double-blind, randomised, placebo-controlled challenge studies have consistently demonstrated that, compared with placebo, caffeine is associated with a greater increase in

anxiety in people with panic or generalised anxiety disorders.²⁵⁷⁻²⁶¹ Similarly, a double-blind, placebo-controlled challenge study found that caffeine increased anxiety levels in people with social phobia.²⁶² Evidence for longer-term effects of caffeine on anxiety is weak. Survey studies have demonstrated a significant association between daily caffeine intake and anxiety among people with panic disorder^{255,256} or high trait anxiety when under stress.²⁶³ However, another study reported no association between daily caffeine intake and anxiety.²⁶⁴ There have been no controlled studies of the long-term effects of caffeine in patients with anxiety disorder.

Conclusion: There is some evidence that caffeine increases anxiety levels and precipitates panic attacks among people with anxiety disorders. However, there is a need for well designed RCTs of the long-term effect on anxiety of reducing or eliminating caffeine intake in people with anxiety disorders.

Carbohydrate-rich, protein-poor diet

Description: A carbohydrate-rich, protein-poor diet has more cereal products and little or no meat, dairy products or nuts.

Rationale: Biochemical imbalances in the brain can mediate or exacerbate the negative effects of stress experienced by individuals. In particular, serotonin activity in the brain occurs as a result of stress. Those with chronic stress may have depleted levels of brain serotonin, which results in their dealing with stress less effectively.²⁶⁵ A diet which increases levels of serotonin in the brain may assist individuals to cope more effectively with stress. Diets high in carbohydrates and low in protein have been reported to increase serotonin levels in the brain.²⁶⁶

Level of evidence: V.

Review of effectiveness: In one RCT, stress-prone individuals on a high-carbohydrate, low-protein diet performed significantly better on memory scanning tasks after experiencing controllable stress compared with those who had a low-carbohydrate, protein-rich diet.²⁶⁵ This difference did not occur, however, when they were subjected to uncontrollable stress. The numbers in this study were very small, just 22 stress-prone subjects and 21 controls, and the stress measured was that experienced in a laboratory setting.

Conclusion: The effects of this diet in people with anxiety disorders have not been rigorously evaluated.

Nicotine avoidance

Description: Nicotine is found in tobacco and in products to assist tobacco withdrawal.

Rationale: Nicotine stimulates the release of various neurotransmitters that can affect anxiety, including nicotinic acetylcholine receptors, glutamate, GABA, dopamine, norepinephrine and serotonin.

Level of evidence: V.

Review of effectiveness: It is now hypothesised that nicotine use results in smokers having increased levels of anxiety, but that nicotine withdrawal also increases anxiety, which is then relieved in the short term by smoking again.²⁶⁷ While abrupt withdrawal of cigarettes can often trigger higher levels of anxiety in the short term,²⁶⁸ these eventually disappear, leaving individuals with fewer symptoms of anxiety than they experienced while regularly smoking.^{269,270} In a study of 70 smokers whose abstinence from smoking was carefully monitored over a 4-week period, West and Hajek reported reductions in anxiety levels.²⁷⁰ However, there are no studies specifically on people with anxiety disorders.

Conclusion: There is currently no evidence on whether abstinence from nicotine benefits people with anxiety disorders.

DISCUSSION

For ease of reference, Box 3 gives an overview of the evidence available for each treatment. The treatments with the best evidence of effectiveness are kava (for generalised anxiety), exercise (for generalised anxiety), relaxation training (for generalised anxiety, panic disorder, dental phobia and test anxiety) and bibliotherapy (for specific phobias). However, none of these treatments has as much support as standard treatments recommended in clinical practice guidelines. For example, a meta-analysis of treatments for generalised anxiety included 13 studies of 22 interventions using cognitive behaviour therapy and 24 studies of 39 interventions using pharmacological treatments.¹³¹ By contrast, for generalised anxiety, we found 11 trials of kava, 11 of exercise, 15 of relaxation and four of bibliotherapy.

More limited evidence supports the effectiveness of acupuncture, music, autogenic training and meditation for generalised anxiety; inositol in the treatment of panic disorder and obsessive-compulsive disorder; massage in the treatment of post-traumatic stress in children; and alcohol avoidance by people with alcohol-use disorders to reduce a range of anxiety disorders. Some of these last treatments may well be effective, but they have received very little research attention. Research on the effectiveness of treatments for anxiety disorders has tended to focus on a small number of standard treatments and needs to be broadened, particularly in view of the public's more favourable attitudes to some non-standard treatments.

Since most of the studies reviewed have dealt with generalised anxiety disorder or high levels of anxiety, there is a particular need for work on other anxiety disorders. Future studies also need to pay particular attention to the use of larger sample sizes, and longer follow-up periods, blinding of treatments (where feasible) and use of intention-to-treat analysis. We also know little about how some of these treatments perform in population subgroups, such as children and adolescents, the elderly, and in perinatal women.

We have previously reviewed the evidence on complementary and self-help treatments for depression.⁶⁰ Anxiety and depression show considerable comorbidity, and prolonged anxiety appears to be an important factor in the aetiology of depression.²⁷¹ Thus, it might be expected that treatments that work for anxiety would also work for depression. However, some treatments that have been reported as effective for anxiety disorders have been largely ignored for depression (eg, kava), and some treatments that have been reported as effective for depression have been largely ignored for anxiety disorders (eg, St John's wort). There is scope for further research extending treatments across the divide between anxiety and depression.

Given the frequent use of complementary and self-help treatments, it would be wise for GPs and others who are treating depressed or anxious patients to routinely enquire about the use of these other treatments. An important reason is to prevent potentially harmful interactions with conventional treatments. There has already been a voluntary recall of kava products and there has been a warning from the Therapeutic Goods Administration about the interaction of St John's wort with prescription medications.

Another reason to enquire about use of complementary and self-help treatments is to educate patients to make better choices. If

3 Evidence on effectiveness of complementary and self-help treatments for anxiety disorders

Treatment	Level of evidence	Conclusion
<i>Medicines and homoeopathic remedies</i>		
Bach flower remedies	II	Not effective for test anxiety. No evidence on other anxiety disorders
Berocca	II	No evidence on anxiety disorders
Ginger	V	No evidence on anxiety disorders
Gotu kola	V	No evidence on anxiety disorders
Combined preparations (including herbs)	II	No convincing evidence
Homoeopathy	II	No convincing evidence
Inositol	II	Some evidence for effectiveness with panic disorder and obsessive-compulsive disorder
Kava	I	Effective for generalised anxiety, but may cause liver toxicity in rare cases
Lemongrass leaves	II	Not effective
Licorice	IV	No convincing evidence
Magnesium	II	No convincing evidence
Passionflower	II	No convincing evidence
St John's wort	IV	No convincing evidence
Valerian	II	No convincing evidence
Vitamin C	V	No convincing evidence
5-hydroxy-L-tryptophan	II	Some evidence for effectiveness, but cannot be recommended because of possible risk of eosinophilia–myalgia syndrome
<i>Physical treatments</i>		
Acupuncture	II	Some evidence for effectiveness for generalised anxiety
Aromatherapy	II	No evidence on anxiety disorders
Hydrotherapy	IV	No convincing evidence
Massage/touch therapy	II	Some evidence for effectiveness with post-traumatic stress disorder in children, but no convincing evidence for adults or on other anxiety disorders
<i>Lifestyle</i>		
Autogenic training	I	Some evidence for effectiveness for generalised anxiety
Bibliotherapy	I	Evidence of effectiveness for specific phobias, but not for other anxiety disorders
Dance/movement therapy	II	Some evidence of effectiveness for test anxiety. No convincing evidence on other anxiety disorders
Exercise	I	Evidence for effectiveness for generalised anxiety. More evidence on other anxiety disorders is needed
Humour	V	No convincing evidence
Meditation	II	Some evidence for effectiveness with generalised anxiety. More evidence on other anxiety disorders is needed
Music	II	Some evidence for effectiveness with generalised anxiety. More evidence on other anxiety disorders is needed
Prayer	II	No convincing evidence
Relaxation	I or II	Evidence for effectiveness with generalised anxiety, panic disorder, dental phobia and test anxiety
Yoga	II	No convincing evidence
<i>Dietary and other changes</i>		
Alcohol	IV	Some evidence that eliminating alcohol reduces anxiety in people with alcohol disorders
Caffeine reduction	II or IV	No convincing evidence
Carbohydrate-rich, protein-poor diet	V	No convincing evidence
Nicotine avoidance	V	No evidence on anxiety disorders

patients do wish to use such treatments, it is preferable that they use those most supported by evidence.

ACKNOWLEDGEMENTS

We wish to thank Tracey Davenport for editorial assistance with this article. Funding was provided by Program Grant No. 179805 from the National Health and Medical Research Council.

COMPETING INTERESTS

None identified.

REFERENCES

- Andrews G, Hall W. The mental health of Australians. Canberra: Mental Health Branch, Australian Government Department of Health and Aged Care, 1999.
- Mathers C, Vos T, Stevenson C. The burden of disease and injury in Australia. Canberra: Australian Institute of Health and Welfare, 1999.
- Goldberg D. Plato versus Aristotle: categorical and dimensional models for common mental disorders. *Compr Psychiatry* 2000; 41 (2 Suppl 1): 8-13.
- National Health Committee. Guidelines for assessing and treating anxiety disorders. Wellington: NHC, 1998.
- Issakidis C, Andrews G. Service utilisation for anxiety in an Australian community sample. *Soc Psychiatry Psychiatr Epidemiol* 2002; 37: 153-163.
- Jorm AF, Medway J, Christensen H, et al. Public beliefs about the helpfulness of interventions for depression: effects on actions taken when experiencing anxiety and depression symptoms. *Aust N Z J Psychiatry* 2000; 34: 619-626.
- MacLennan AH, Wilson DH, Taylor AW. The escalating cost and prevalence of alternative medicine. *Prev Med* 2002; 35: 166-173.
- Jorm AF, Christensen H, Griffiths KM, Rodgers B. Effectiveness of complementary and self-help treatments for depression. *Med J Aust* 2002; 176 (10 Suppl): S84-S96.
- American Academy of Child and Adolescent Psychiatry. Practice parameters for the assessment and treatment of children and adolescents with obsessive-compulsive disorder. *J Am Acad Child Adolesc Psychiatry* 1998; 37 (10 Suppl): 27S-45S.
- American Academy of Child and Adolescent Psychiatry. Practice parameters for the assessment and treatment of children and adolescents with posttraumatic stress disorder. *J Am Acad Child Adolesc Psychiatry* 1998; 37 (10 Suppl): 4S-26S.
- American Psychiatric Association. Practice guideline for the treatment of patients with panic disorder. Work group on panic disorder. *Am J Psychiatry* 1998; 155 (5 Suppl): 1-34.
- Bernstein GA, Shaw K. Practice parameters for the assessment and treatment of children and adolescents with anxiety disorders. American Academy of Child and Adolescent Psychiatry. *J Am Acad Child Adolesc Psychiatry* 1997; 36 (10 Suppl): 69S-84S.
- Department of Health. Treatment choice in psychological therapies and counselling — evidence based clinical practice guideline. London: Department of Health, 2001.
- Foa EB, Davidson JRT, Frances A. The expert consensus guideline series. Treatment of posttraumatic stress disorder. *J Clin Psychiatry* 1999; 60 Suppl 16: 3-76.
- Foa EB, Keane TM, Friedman MJ. Guidelines for treatment of PTSD. *J Trauma Stress* 2000; 13: 539-588.
- Institute for Clinical Systems Improvement. Health care guideline: major depression, panic disorder and generalised anxiety disorder in adults in primary care. Bloomington: ICSI, 2001.
- National Department of Health. Standard treatment guidelines and essential drugs list for South Africa: adult hospital level. Pretoria: South African Department of Health, 1998.
- Royal Australian and New Zealand College of Psychiatrists Clinical Practice Guidelines Team for Panic Disorder and Agoraphobia. Australian and New Zealand clinical practice guidelines for treatment of panic disorder and agoraphobia. *Aust N Z J Psychiatry* 2003; 37: 641-656.
- World Health Organization. WHO guide to mental health in primary care. London: Royal Society of Medicine, 2000.
- Andrews G, Crino R, Hunt C, et al. The treatment of anxiety disorders: clinician's guide and patient manuals. 1st ed. New York: Cambridge University Press, 1994.
- Fugh-Berman A, Cott JM. Dietary supplements and natural products as psychotherapeutic agents. *Psychosom Med* 1999; 61: 712-728.
- Muskin PR, editor. Complementary and alternative medicine and psychiatry. Washington, DC: American Psychiatric Association, 2000.
- Nathan PE, Gorman JM, editors. A guide to treatments that work. New York: Oxford University Press, 1998.
- Wong AH, Smith M, Boon HS. Herbal remedies in psychiatric practice. *Arch Gen Psychiatry* 1998; 55: 1033-1044.
- National Health and Medical Research Council. How to use the evidence: assessment and application of scientific evidence. Canberra: NHMRC, 2000.
- Armstrong NC, Ernst E. A randomized, double-blind placebo-controlled trial of a Bach Flower remedy. *Complement Ther Nurs Midwifery* 2001; 7: 215-221.
- Walach H, Rilling C, Engelke U. Efficacy of Bach-Flower remedies in test anxiety: a double-blind, placebo-controlled, randomized trial with partial crossover. *J Anxiety Disord* 2001; 15: 359-366.
- Carroll D, Ring C, Suter M, Willemsen G. The effects of an oral multivitamin combination with calcium, magnesium, and zinc on psychological well-being in healthy young male volunteers: a double-blind placebo-controlled trial. *Psychopharmacologia* 2000; 150: 220-225.
- Topic B, Hasenohrl RU, Hacker R, Huston JP. Enhanced conditioned inhibitory avoidance by a combined extract of *Zingiber officinale* and *Ginkgo biloba*. *Phytother Res* 2002; 16: 312-315.
- Mantani R, Cimino A. A primer of complementary and alternative medicine and its relevance in the treatment of mental health problems. *Psychiatr Q* 2002; 73: 367-381.
- Miller LG. Herbal medicinals: selected clinical considerations focusing on known or potential drug-herb interactions. *Arch Intern Med* 1998; 158: 2200-2211.
- Vaes LP, Chyka PA. Interactions of warfarin with garlic, ginger, ginkgo or ginseng: nature of the evidence. *Ann Pharmacother* 2000; 34: 1478-1482.
- Bradwejn JC, Zhou Y, Koszycki D, Shlik JA. Double-blind, placebo-controlled study on the effects of Gotu Kola (*Centella asiatica*) on acoustic startle response in healthy subjects. *J Clin Psychopharmacol* 2000; 20: 680-684.
- Chen HC, Hsieh MT, Lai E. Studies on the suanzaorentang in the treatment of anxiety. *Psychopharmacology (Berl)* 1985; 85: 486-487.
- Mills PJ, Farag NH, Newton RP, Parry BL. Effects of a traditional herbal supplement on anxiety in patients with generalized anxiety disorder. *J Clin Psychopharmacol* 2002; 22: 443-444.
- Panijel M. Treatment of moderately severe anxiety states. *Therapiewoche* 1985; 35: 4659-4668.
- Bourin M, Bougerol T, Guitton B, Broutin E. A combination of plant extracts in the treatment of outpatients with adjustment disorder with anxious mood: controlled study versus placebo. *Fundam Clin Pharmacol* 1997; 11: 127-132.
- Slonim D, White K. Homeopathy and psychiatry. *J Mind Behav* 1983; 4: 401-410.
- Fisher P. How does homeopathy work: are we looking in the right place? *Homeopathy* 2003; 92: 1-2.
- Cialdella P, Boissel JP, Belon P. Homeopathic specialities as substitutes for benzodiazepines: double-blind vs placebo study. *Therapie* 2001; 56: 397-402.
- McCutcheon LE. Treatment of anxiety with a homeopathic remedy. *J Applied Nutr* 1996; 48: 2-6.
- Benjamin J, Agam G, Levine J, et al. Inositol treatment in psychiatry. *Psychopharmacol Bull* 1995; 31: 167-175.
- Benjamin J, Levine J, Fux M, et al. Double-blind, placebo-controlled, crossover trial of inositol treatment for panic disorder. *Am J Psychiatry* 1995; 152: 1084-1086.
- Palatnik A, Frolov K, Fux M, Benjamin J. Double-blind, controlled, crossover trial of inositol versus fluvoxamine for the treatment of panic disorder. *J Clin Psychopharmacol* 2001; 21: 335-339.
- Fux M, Levine J, Aviv A, Belmaker RH. Inositol treatment of obsessive-compulsive disorder. *Am J Psychiatry* 1996; 153: 1219-1221.
- Fux M, Benjamin J, Belmaker RH. Inositol versus placebo augmentation of serotonin reuptake inhibitor in the treatment of obsessive-compulsive disorder: a double-blind cross-over study. *Int J Neuropsychopharmacol* 1999; 2: 193-195.

- 47 Kaplan Z, Amir M, Swartz M, Levine J. Inositol treatment of post-traumatic stress disorder. *Anxiety* 1996; 2: 51-52.
- 48 Cairney S, Maruff P, Clough AR. The neurobehavioural effects of kava. *Aust N Z J Psychiatry* 2002; 36: 657-662.
- 49 Pittler MH, Ernst E. Kava extract for treating anxiety [Cochrane Review]. The Cochrane Library, Issue 1, 2003.
- 50 Bilia AR, Gallon S, Vincieri FF. Kava-kava and anxiety: growing knowledge about the efficacy and safety. *Life Sci* 2002; 70: 2581-2597.
- 51 Leite JR, Seabra M de L, Maluf E, et al. Pharmacology of lemongrass (*Cymbopogon citratus Stapf*). III. Assessment of eventual toxic, hypnotic and anxiolytic effects on humans. *J Ethnopharmacol* 1986; 17: 75-83.
- 52 Duke JA. Novel psychotherapeutic drugs: a role for ethnobotany: commentary. *Psychopharmacol Bull* 1995; 31: 177-184.
- 53 Simon W, Edwards RV. Glycyrrhiza (licorice) in the treatment of psychiatric illness. *J Clin Exp Psychopathol* 1957; 18: 79-86.
- 54 Olukoga A, Donaldson D. Licorice and its health implications. *J R Soc Health* 2000; 120: 83-89.
- 55 Strandberg TE, Anderssen S, Jarvenpaa AL, McKeigue PM. Preterm birth and licorice consumption during pregnancy. *Am J Epidemiol* 2002; 156: 803-805.
- 56 Kirov GK, Tsachev KN. Magnesium, schizophrenia and manic-depressive disease. *Neuropsychobiology* 1990; 23: 79-81.
- 57 Bockova E, Hronek J, Kolomaznik M, et al. Potentiation of the effects of anxiolytics with magnesium salts. *Cesk Psychiatr* 1992; 88: 141-144.
- 58 Bilia R, Bergonzi MC, Gallori S, et al. Stability of the constituents of Calendula, milk thistle and passionflower tinctures by LC-DAD and LC-MS. *J Pharm Biomed Anal* 2002; 30: 613-624.
- 59 Akhondzadeh S, Naghavi HR, Vazirian M, et al. Passionflower in the treatment of generalised anxiety: a pilot double-blind randomised controlled trial with oxazepam. *J Clin Pharm Ther* 2001; 26: 363-367.
- 60 Nathan P. The experimental and clinical pharmacology of St John's Wort (*Hypericum perforatum* L.). *Mol Psychiatry* 1999; 4: 333-338.
- 61 Taylor LH, Kobak KA. An open-label trial of St. John's Wort (*Hypericum perforatum*) in obsessive-compulsive disorder. *J Clin Psychiatry* 2000; 61: 575-578.
- 62 Davidson JR, Connor KM. St. John's Wort in generalized anxiety disorder: three case reports. *J Clin Psychopharmacol* 2001; 21: 635-636.
- 63 Dominguez RA. Valerian: its value as a sedative hypnotic. In: Mischoulon D, Rosenbaum JF, editors. *Natural medications for psychiatric disorders: considering the alternatives*. 1st ed. Philadelphia: Lippincott Williams and Wilkins, 2002: 132-146.
- 64 Andreatini R, Sartori VA, Seabra ML, Leite JR. Effect of valepotriates (valerian extract) in generalized anxiety disorder: a randomized placebo-controlled pilot study. *Phytother Res* 2002; 16: 650-654.
- 65 Kohlen R, Oswald D. The effects of valerian, propranolol, and their combination on activation, performance, and mood of healthy volunteers under social stress conditions. *Pharmacopsychiatry* 1988; 21: 447-448.
- 66 Willey LB, Mady SP, Cobaugh DJ, Wax PM. Valerian overdose: a case report. *Vet Hum Toxicol* 1995; 37: 364-365.
- 67 Yager J, Siegfried SL, DiMatteo TL. Use of alternative remedies by psychiatric patients: illustrative vignettes and a discussion of the issues. *Am J Psychiatry* 1999; 156: 1432-1438.
- 68 Wong AH, Smith M, Boon HS. Herbal remedies in psychiatric practice. *Arch Gen Psychiatry* 1998; 55: 1033-1044.
- 69 Mischoulon D, Rosenbaum JF, editors. *Natural medications for psychiatric disorders: considering the alternatives*. 1st ed. Philadelphia: Lippincott Williams and Wilkins, 2002.
- 70 Reifenberg E. On a case of obsessive-compulsive neurosis significantly improved with ascorbic acid. *Psychiatr Neurol Med Psychol (Leipz)* 1954; 5: 161-167.
- 71 Westenberg HG, den Boer JA, Kahn RS. Psychopharmacology of anxiety disorders: on the role of serotonin in the treatment of anxiety states and phobic disorders. *Psychopharmacol Bull* 1987; 23: 145-149.
- 72 Kahn RS, Westenberg HG, Verhoeven WM, et al. Effect of a serotonin precursor and uptake inhibitor in anxiety disorders: a double-blind comparison of 5-hydroxytryptophan, clomipramine and placebo. *Int Clin Psychopharmacol* 1987; 2: 33-45.
- 73 Schruers K, Pols H, Overbeek T, et al. 5-hydroxytryptophan inhibits 35% CO₂ induced panic. *Int J Neuropsychopharmacol* 2000; 3 Suppl 1: 272.
- 74 Kahn RS, Westenberg HG. L-5-Hydroxytryptophan in the treatment of anxiety disorders. *J Affect Disord* 1985; 8: 197-200.
- 75 Blier P, Bergeron R. Sequential administration of augmentation strategies in treatment-resistant obsessive-compulsive disorder: preliminary findings. *Int Clin Psychopharmacol* 1996; 11: 37-44.
- 76 Larzelere MM, Wiseman P. Anxiety, depression, and insomnia. *Prim Care* 2002; 29: 339-360.
- 77 Cauffield JS, Forbes HJ. Dietary supplements used in the treatment of depression, anxiety, and sleep disorders. *Lippincott's Prim Care Pract* 1999; 3: 290-304.
- 78 US Food and Drug Administration. Information paper on L-tryptophan and 5-hydroxy-L-tryptophan, February 2001. Available at: vm.cfsan.fda.gov/~dms/ds-tryp1.html (accessed Mar 2004).
- 79 Guizhen L, Yunjun Z, Linxiang G, Aizhen L. Comparative study on acupuncture combined with behavioral desensitization for treatment of anxiety neuroses. *Am J Acupunct* 1998; 26: 117-120.
- 80 Spector IP, Carey MP, Jorgensen RS, et al. Cue-controlled relaxation and "aromatherapy" in the treatment of speech anxiety. *Behav Cognit Psychother* 1993; 21: 239-253.
- 81 Cooke B, Ernst E. Aromatherapy: a systematic review. *Br J Gen Pract* 2000; 50: 493-496.
- 82 Levine BA. Use of hydrotherapy in reduction of anxiety. *Psychol Rep* 1984; 55: 526.
- 83 Field TM. Touch therapies. In: Hoffman RR, Sherrick MF, Warm JS, editors. *Viewing psychology as a whole: the integrative science of William N Dember*. 1st ed. New York: American Psychological Association, 1998: 603-624.
- 84 Vickers A, Zollman C. ABC of complementary medicine: massage therapies. *BMJ* 1999; 319: 1254-1257.
- 85 Field T, Seligman S, Scafidi F, Schanberg S. Alleviating posttraumatic stress in children following Hurricane Andrew. *J Appl Dev Psychol* 1996; 17: 37-50.
- 86 Field T, Grizzle N, Scafidi F, Schanberg S. Massage and relaxation therapies' effects on depressed adolescent mothers. *Adolescence* 1996; 31: 903-911.
- 87 Field T, Ironson G, Scafidi F, et al. Massage therapy reduces anxiety and enhances EEG pattern of alertness and math computations. *Int J Neurosci* 1996; 86: 197-205.
- 88 Shulman KR, Jones GE. The effectiveness of massage therapy intervention on reducing anxiety in the workplace. *J Appl Behav Sci* 1996; 32: 160-173.
- 89 Richards KC, Gibson R, Overton-McCoy AL. Effects of massage in acute and critical care. *AACN Clin Issues* 2000; 11: 77-96.
- 90 Stetter F, Kupper S. Autogenic training: a meta-analysis of clinical outcome studies. *Appl Psychophysiol Biofeedback* 2002; 27: 45-98.
- 91 Scogin F, Bynum J, Stephens G, Calhoun S. Efficacy of self-administered treatment programs: meta-analytic review. *Prof Psychol - Res Pract* 1990; 21: 42-47.
- 92 Gould RA, Clum GA, Shapiro D. The use of bibliotherapy in the treatment of panic: a preliminary investigation. *Behav Ther* 1993; 24: 241-252.
- 93 Marrs RW. A meta-analysis of bibliotherapy studies. *Am J Community Psychol* 1995; 23: 843-870.
- 94 Newman MG, Erickson T, Przeworski A, Dzus E. Self-help and minimal-contact therapies for anxiety disorders: is human contact necessary for therapeutic effect? *J Clin Psychol* 2003; 59: 271-274.
- 95 Ritter M, Low KG. Effects of dance/movement therapy: a meta-analysis. *The Arts in Psychotherapy* 1996; 23: 249-260.
- 96 Kline F, Burgoyne RW, Staples F, et al. A report on the use of movement therapy for chronic, severely disabled outpatients. *Art Psychother* 1977; 4: 181-183.
- 97 Brooks D, Stark A. The effect of dance/movement therapy on affect: a pilot study. *Am J Dance Ther* 1989; 11: 101-112.
- 98 Erwin-Grabner T, Goodill SW, Hill ES, Von Neida K. Effectiveness of dance/movement therapy on reducing test anxiety. *Am J Dance Ther* 1999; 21: 19-34.
- 99 Kubota N. An analysis of the therapeutic process of Dosa therapy. *Dissertation Abstracts International* 2001; 61: 3352.
- 100 Kuettel TJ. Affective change in dance therapy. *Am J Dance Ther* 1982; 5: 56-64.
- 101 Leste A, Rust J. Effects of dance on anxiety. *Percept Mot Skills* 1984; 58: 767-772.
- 102 Tkachuk GA, Martin GL. Exercise therapy for patients with psychiatric disorders: research and clinical implications. *Prof Psychol - Res Pract* 1999; 30: 275-282.

- 103 Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exerc* 2001; 33 Suppl: 587-597.
- 104 Petruzzello SJ, Landers DM, Hatfield BD, et al. A meta-analysis on the anxiety-reducing effects of acute and chronic exercise. Outcomes and mechanisms. *Sports Med* 1991; 11: 143-182.
- 105 Stich FA. A meta-analysis of physical exercise as a treatment for symptoms of anxiety and depression [dissertation]. Madison: University of Wisconsin, 1998.
- 106 Steptoe A, Edwards S, Moses J, Mathews A. The effects of exercise training on mood and perceived coping ability in anxious adults from the general population. *J Psychosom Res* 1989; 33: 537-547.
- 107 McEntee DJ, Halgin RP. Cognitive group therapy and aerobic exercise in the treatment of anxiety. *J Coll Student Psychother* 1999; 13: 37-55.
- 108 Broocks A, Bandelow B, Pekrun G, et al. Comparison of aerobic exercise, clomipramine, and placebo in the treatment of panic disorder. *Am J Psychiatry* 1998; 155: 603-609.
- 109 Martinsen EW, Hoffart A, Solberg OY. Aerobic and non-aerobic forms of exercise in the treatment of anxiety disorders. *Stress Med* 1989; 5: 115-120.
- 110 Manger TA. The effects of a community-based aerobic exercise program on posttraumatic stress disorder-related symptoms and concomitant anxiety and depression [dissertation]. Hempstead, NY: Hofstra University, 2000.
- 111 Ventis WL, Higbee G, Murdock SA. Using humor in systematic desensitization to reduce fear. *J Gen Psychol* 2001; 128: 241-253.
- 112 Abel MH. Interaction of humor and gender in moderating relationships between stress and outcomes. *J Psychol* 1998; 132: 267-276.
- 113 Yovetich NA, Dale JA, Hudak MA. Benefits of humor in reduction of threat-induced anxiety. *Psychol Rep* 1990; 66: 51-58.
- 114 Ascough JC, Ettinger RF, Nelson DA. Humour, anxiety and task performance. *J Pers Soc Psychol* 1971; 19: 243-246.
- 115 Thomas D, Abbas KA. Comparison of transcendental meditation and progressive relaxation in reducing anxiety. *BMJ* 1978; 2: 1749.
- 116 Lehrer PM, Woolfolk RL, Rooney AJ, et al. Progressive relaxation and meditation. A study of psychophysiological and therapeutic differences between two techniques. *Behav Res Ther* 1983; 21: 651-662.
- 117 Raskin M, Bali LR, Peeke HV. Muscle biofeedback and transcendental meditation. A controlled evaluation of efficacy in the treatment of chronic anxiety. *Arch Gen Psychiatry* 1980; 37: 93-97.
- 118 Wilson HB. The specific effects model: relaxation and meditation effects on cognitive and somatic anxiety [dissertation]. Athens, Ohio: Ohio University, 2000.
- 119 Boswell PC, Murray EJ. Effects of meditation on psychological and physiological measures of anxiety. *J Consult Clin Psychol* 1979; 47: 606-607.
- 120 Kirkland K, Hollandsworth JG. Effective test taking: skills-acquisition versus anxiety-reduction techniques. *J Consult Clin Psychol* 1980; 48: 431-439.
- 121 Shannahoff-Khalsa DS, Ray LE, Levine S, et al. Randomized controlled trial of yogic meditation techniques for patients with obsessive-compulsive disorder. *CNS Spectr* 1999; 4: 34-47.
- 122 Morse DR. An exploratory study of the use of meditation alone and in combination with hypnosis in clinical dentistry. *J Am Soc Psychosomat Dent Med* 1977; 24: 113-120.
- 123 Morse DR, Cohen BB. Desensitization using meditation-hypnosis to control "needle" phobia in two dental patients. *Anesth Prog* 1983; 30: 83-85.
- 124 Russell LA. Comparisons of cognitive, music, and imagery techniques on anxiety reduction with university students. *J Coll Student Dev* 1992; 33: 516-523.
- 125 Stanton HE. A comparison of the effects of an hypnotic procedure and music on anxiety level. *Aust J Clin Exp Hypn* 1984; 12: 127-132.
- 126 Wu SM. Effects of music therapy on anxiety, depression and self-esteem of undergraduates. *Psychologica* 2002; 45: 104-114.
- 127 Stoudenmire J. A comparison of muscle relaxation training and music in the reduction of state and trait anxiety. *J Clin Psychol* 1975; 31: 490-492.
- 128 Kerr T, Walsh J, Marshall A. Emotional change processes in music-assisted reframing. *J Music Ther* 2001; 38: 193-211.
- 129 Razali SM, Hasanah CI, Aminah K, Subramaniam M. Religious-sociocultural psychotherapy in patients with anxiety and depression. *Aust N Z J Psychiatry* 1998; 32: 867-872.
- 130 Snaith P. Meditation and psychotherapy. *Br J Psychiatry* 1998; 173: 193-195.
- 131 Gould RA, Otto MW, Pollack MH, Yap L. Cognitive behavioral and pharmacological treatment of generalized anxiety disorder: a preliminary meta-analysis. *Behav Ther* 1997; 28: 285-305.
- 132 Boswell PC, Murray EJ. Effects of meditation on psychological and physiological measures of anxiety. *J Consult Clin Psychol* 1979; 47: 606-607.
- 133 Lehrer PM, Atthowe JM, Webber ESP. Effects of progressive relaxation and autogenic training on anxiety and physiological measures, with some data on hypnotizability. In: McGuigan FJ, Sime WE, Wallace JM, editors. *Stress and tension control*. New York: Plenum, 1980: 171-181.
- 134 Hillenberg JB, Collins FL. The importance of home practice for progressive relaxation training. *Behav Res Ther* 1983; 21: 633-642.
- 135 Flanders P, McNamara JR. Relaxation training and home practice in the treatment of anxiety. *Psychol Rep* 1987; 61: 819-822.
- 136 Cragan MK, Deffenbacher JL. Anxiety management training and relaxation as self-control in the treatment of generalized anxiety in medical outpatients. *J Counseling Psychol* 1984; 31: 123-131.
- 137 DeBerry S, Davis S, Reinhard KE. A comparison of meditation-relaxation and cognitive/behavioural techniques for reducing anxiety and depression in a geriatric hospital. *J Geriatr Psychiatry* 1989; 22: 231-247.
- 138 Tarrier N, Main CJ. Applied relaxation training for generalised anxiety and panic attacks: the efficacy of a learnt coping strategy on subjective reports. *Br J Psychiatry* 1986; 149: 330-336.
- 139 Carlson CR, Collins FL Jr, Nitz AJ, et al. Muscle stretching as an alternative relaxation training procedure. *J Behav Ther Exp Psychiatry* 1990; 21: 29-38.
- 140 O'Neill LM, Barnier AJ, McConkey K. Treating anxiety with self-hypnosis and relaxation. *Contemp Hypn* 1999; 16: 68-80.
- 141 Barlow DH, Rapee RM, Brown TA. Behavioural treatment of generalised anxiety disorder. *Behav Ther* 1992; 23: 551-570.
- 142 Borkovec TD, Costello E. Efficacy of applied relaxation and cognitive-behavioural therapy in the treatment of generalised anxiety disorder. *J Consult Clin Psychol* 1993; 61: 611-619.
- 143 Ost LG, Breitholtz E. Applied relaxation vs. cognitive therapy in the treatment of generalized anxiety disorder. *Behav Res Ther* 2000; 38: 777-790.
- 144 Kumaraiah V. Treatment of anxiety through GSR biofeedback and progressive muscular relaxation. *Indian J Clin Psychol* 1979; 6: 209-211.
- 145 Hutchings DF, Denney DR, Basgall J, Houston BK. Anxiety management and applied relaxation in reducing general anxiety. *Behav Res Ther* 1980; 18: 181-190.
- 146 LeBoeuf A, Lodge JA. A comparison of frontalis EMG feedback training and progressive relaxation in the treatment of chronic anxiety. *Br J Psychiatry* 1980; 137: 279-284.
- 147 Rupert PA, Dobbins K, Mathew RJ. EMG biofeedback and relaxation instructions in the treatment of chronic anxiety. *Am J Clin Biofeedback* 1981; 4: 52-61.
- 148 Pitkin GL. An investigation of cognitive and relaxation-imagery techniques on college students' level of general anxiety [dissertation]. Pullman: Washington State University, 1983.
- 149 Bouchard SJ. Effects of a self-administered subliminal-relaxation treatment on anxiety [dissertation]. San Diego, Calif: United States International University, 1984.
- 150 DeBerry S. An evaluation of progressive muscle relaxation on stress related symptoms in a geriatric population. *Int J Aging Hum Dev* 1981; 14: 255-269.
- 151 Walker LS, Busby K, Walker JL. The differential effects of progressive relaxation and imagery on anxiety. *Can J Counselling* 1987; 21: 207-215.
- 152 Rowe GL. The effects of matching individual anxiety response patterns to treatment [dissertation]. Pullman: Washington State University, 1991.
- 153 Kohli A, Varma VK, Nehra R. Comparison of efficacy of psychorelaxation and pharmacotherapy in generalized anxiety disorder. *J Pers Clin Stud* 2000; 16: 43-48.
- 154 Gould RA, Otto MW, Pollack MH. A meta-analysis of treatment outcome for panic disorder. *Clin Psychol Rev* 1995; 15: 819-844.
- 155 Ost LG, Jerremalm A, Jansson L. Individual response patterns and the effects of different behavioural methods in the treatment of agoraphobia. *Behav Res Ther* 1984; 22: 697-707.
- 156 Ost LG. Applied relaxation: description of a coping technique and review of controlled studies. *Behav Res Ther* 1987; 25: 397-409.
- 157 McNamee G, O'Sullivan G, Lelliott PT, Marks I. Telephone-guided treatment for housebound agoraphobics with panic disorder: exposure vs. relaxation. *Behav Ther* 1989; 20: 491-497.

- 158 Michelson L, Mavissakalian M, Marchione K. Cognitive, behavioural and psychophysiological treatments of agoraphobia: a comparative outcome investigation. *Behav Ther* 1988; 19: 97-120.
- 159 Barlow DH, Craske MG, Cerny JA, Klosko JS. Behavioural treatment of panic disorder. *Behav Ther* 1989; 20: 261-282.
- 160 Ost LG, Westling BE, Hellstron K. Applied relaxation, exposure in vivo and cognitive methods in the treatment of panic disorder with agoraphobia. *Behav Res Ther* 1993; 31: 383-394.
- 161 Marks IM, Swinson RP, Basoglu M, et al. Alprazolam and exposure alone and combined in panic disorder with agoraphobia. A controlled study in London and Toronto. *Br J Psychiatry* 1993; 162: 776-787.
- 162 Clark DM, Salkovskis PM, Hackmann A, et al. A comparison of cognitive therapy, applied relaxation and imipramine in the treatment of panic disorder. *Br J Psychiatry* 1994; 164: 759-769.
- 163 Beck JG, Stanley MA, Baldwin LE, et al. A comparison of cognitive therapy and relaxation training for panic disorder. *J Consult Clin Psychol* 1994; 62: 818-826.
- 164 Ost LG, Westling BE. Applied relaxation vs. cognitive behaviour therapy in the treatment of panic disorder. *Behav Res Ther* 1995; 33: 145-158.
- 165 Arntz A, van den Hout M. Cognitive therapy vs. applied relaxation. *Behav Res Ther* 1996; 34: 113-121.
- 166 Van Etten ML, Taylor S. Comparative efficacy of treatments for PTSD: a meta-analysis. *Clin Psychol Psychother* 1998; 5: 126-145.
- 167 Silver SM, Brooks A, Obenchain J. Treatment of Vietnam War veterans with PTSD: a comparison of eye movement desensitization and reprocessing, biofeedback, and relaxation training. *J Trauma Stress* 1995; 8: 337-342.
- 168 Vaughan K, Armstrong MS, Gold R, et al. A trial of eye movement desensitization compared with image habituation training and applied muscle relaxation in post-traumatic stress disorder. *J Behav Ther Exp Psychiatry* 1994; 25: 283-291.
- 169 Echeburua E, de Corral P, Sarasua B, et al. Treatment of acute posttraumatic stress disorder in rape victims: an experimental study. *J Anxiety Disord* 1996; 10: 185-199.
- 170 Watson CG, Tuorila JR, Vickers KS, et al. The efficacies of three relaxation regimens in the treatment of PTSD in Vietnam war veterans. *J Clin Psychol* 1997; 53: 917-923.
- 171 Marks I, Lovell K, Noshirvani H, et al. Treatment of posttraumatic stress disorder by exposure and/or cognitive restructuring: a controlled study. *Arch Gen Psychiatry* 1998; 55: 317-325.
- 172 Taylor S, Thordarson DS, Maxfield L, et al. Comparative efficacy, speed and adverse effects of three PTSD treatments: exposure therapy, EMDR, and relaxation training. *J Consult Clin Psychol* 2003; 71: 330-338.
- 173 Hatton AY. Anxiety sensitivity, perfectionism and risk-taking: a therapeutic intervention for obsessive-compulsive disorder [dissertation]. Columbia: University of South Carolina, 1999.
- 174 Fals-Stewart W, Marks AP, Schafer J. A comparison of behavioral group therapy and individual therapy in treating obsessive-compulsive disorder. *J Nerv Ment Dis* 1993; 181: 189-193.
- 175 Marks IM, Hodgson R, Rachman S. Treatment of chronic obsessive-compulsive neurosis by in-vivo exposure. A two-year follow-up and issues in treatment. *Br J Psychiatry* 1975; 127: 349-364.
- 176 Greist JH, Marks IM, Baer L, et al. Behavior therapy for obsessive-compulsive disorder guided by a computer or by a clinician compared with relaxation as a control. *J Clin Psychiatry* 2002; 63: 138-145.
- 177 Ost LG, Jerremalm A, Johansson J. Individual response patterns and the effects of different behavioural methods in the treatment of social phobia. *Behav Res Ther* 1981; 19: 1-16.
- 178 Jerremalm A, Jansson L, Ost LG. Cognitive and physiological reactivity and the effects of different behavioral methods in the treatment of social phobia. *Behav Res Ther* 1986; 24: 171-180.
- 179 Fedoroff IC, Taylor S. Psychological and pharmacological treatments of social phobia: a meta-analysis. *J Clin Psychopharmacol* 2001; 21: 311-324.
- 180 Miller MP, Murphy PJ, Miller TP. Comparison of electromyographic feedback and progressive relaxation training in treating circumscribed anxiety stress reactions. *J Consult Clin Psychol* 1978; 46: 1291-1298.
- 181 Jerremalm A, Jansson L, Ost LG. Individual response patterns and the effects of different behavioural methods in the treatment of dental phobia. *Behav Res Ther* 1986; 24: 587-596.
- 182 Vassend O, Willumsen T, Hoffart A. Effects of dental fear treatment on general distress. The role of personality variables and treatment method. *Behav Modif* 2000; 24: 580-599.
- 183 Berggren U, Hakeberg M, Carlsson SG. Relaxation vs. cognitively oriented therapies for dental fear. *J Dent Res* 2000; 79: 1645-1651.
- 184 Willumsen T, Vassend O, Hoffart A. A comparison of cognitive therapy, applied relaxation, and nitrous oxide sedation in the treatment of dental fear. *Acta Odontol Scand* 2001; 59: 290-296.
- 185 Bedell JR, Archer RP, Rosmann M. Relaxation therapy, desensitization, and the treatment of anxiety-based disorders. *J Clin Psychol* 1979; 35: 840-843.
- 186 McGlynn FD, Williamson LM, Davis DJ. Semi-automated desensitization as a treatment for genuinely fearful subjects. *Behav Res Ther* 1973; 11: 313-315.
- 187 Beiman I, Israel E, Johnson SA. During training and posttraining effects of live and taped extended progressive relaxation, self-relaxation, and electromyogram biofeedback. *J Consult Clin Psychol* 1978; 46: 314-321.
- 188 Norton GR, Johnson WE. A comparison of two relaxation procedures for reducing cognitive and somatic anxiety. *J Behav Ther Exp Psychiatry* 1983; 14: 209-214.
- 189 Rachman S. Studies in desensitization: I. The separate effects of relaxation and desensitization. *Behav Res Ther* 1965; 3: 245-251.
- 190 Lanyon RI, Manosevitz M, Imber RR. Systematic desensitization: distribution of practice and symptom substitution. *Behav Res Ther* 1968; 6: 323-329.
- 191 Gilroy LJ, Kirkby KC, Daniels BA, et al. Controlled comparison of computer-aided vicarious exposure versus live exposure in the treatment of spider phobia. *Behav Ther* 2000; 31: 733-744.
- 192 Ost LG, Johansson J, Jerremalm A. Individual response patterns and the effects of different behavioural methods in the treatment of claustrophobia. *Behav Res Ther* 1982; 20: 445-460.
- 193 Ost LG, Lindahl IL, Sterner U, Jerremalm A. Exposure in vivo vs applied relaxation in the treatment of blood phobia. *Behav Res Ther* 1984; 22: 205-216.
- 194 Ost LG, Sterner U, Fellenius J. Applied tension, applied relaxation, and the combination in the treatment of blood phobia. *Behav Res Ther* 1989; 27: 109-121.
- 195 Gatchel RJ, Hatch JP, Watson PJ, et al. Comparative effectiveness of voluntary heart rate control and muscular relaxation as active coping skills for reducing speech anxiety. *J Consult Clin Psychol* 1977; 45: 1093-1100.
- 196 Russell RK. The use of systematic desensitization and conditioned relaxation in the treatment of public speaking anxiety [dissertation]. Chicago: University of Illinois, 1972.
- 197 Trexler LD, Karst TO. Rational-emotive therapy, placebo, and no-treatment effects on public-speaking anxiety. *J Abnorm Psychol* 1972; 79: 60-67.
- 198 Gross RT, Fremouw WJ. Cognitive restructuring and progressive relaxation for treatment of empirical subtypes of speech-anxious subjects. *Cognit Ther Res* 1982; 6: 429-436.
- 199 Goldfried MR, Trier CS. Effectiveness of relaxation as an active coping skill. *J Abnorm Psychol* 1974; 83: 348-355.
- 200 Spector IP, Carey MP, Jorgensen RS, et al. Cue-controlled relaxation and "aromatherapy" in the treatment of speech anxiety. *Behav Psychother* 1993; 21: 239-253.
- 201 Osberg JW. The effectiveness of applied relaxation in the treatment of speech anxiety. *Behav Ther* 1981; 12: 723-729.
- 202 Plotkin DP. The effects of interoceptive exposure on fear reduction and return of fear in individuals with public speaking anxiety [dissertation]. Los Angeles: University of California, 2001.
- 203 Laxer RM, Quarter J, Kooman A, Walker K. Systematic desensitization and relaxation of high-test-anxious secondary school students. *J Counseling Psychol* 1969; 16: 446-451.
- 204 Laxer RM, Walker K. Counterconditioning versus relaxation in the desensitization of test anxiety. *J Counseling Psychol* 1970; 17: 431-436.
- 205 Russell RK, Wise F, Stratoudakis JP. Treatment of test anxiety by cue-controlled relaxation and systematic desensitization. *J Counseling Psychol* 1976; 23: 563-566.
- 206 Chang-Liang R, Denney DR. Applied relaxation as training in self-control. *J Counseling Psychol* 1976; 23: 183-189.
- 207 McGlynn FD, Kinjo K, Doherty G. Effects of cue-controlled relaxation, a placebo treatment, and no treatment on changes in self-reported test anxiety among college students. *J Clin Psychol* 1978; 34: 707-714.
- 208 Baither RC, Godsey R. Rational emotive education and relaxation training in large group treatment of test anxiety. *Psychol Rep* 1979; 45: 326.

- 209 Hahnloser RM. A comparison of cognitive restructuring and progressive relaxation in test anxiety reduction [dissertation]. Eugene: University of Oregon, 1974.
- 210 Denney DR. Active, passive, and vicarious desensitization. *J Counseling Psychol* 1974; 21: 369-375.
- 211 Katz B. Treatment of test anxious students by rational-emotive therapy, relaxation placebo and no-treatment [dissertation]. Salt Lake City: University of Utah, 1977.
- 212 Jenkins SA. The relative efficacy of two treatment procedures in modifying maladaptive response systems in the high test anxious student [dissertation]. Bloomington: Indiana University, 1977.
- 213 Johnson DL. The relative efficacy of cue-controlled relaxation therapy, pseudotherapy, and no-treatment control on psychometric and psychophysiological measures of test anxiety [dissertation]. Blacksburg: Virginia Polytechnic Institute and State University, 1977.
- 214 Schafer NB. Differential treatment of test anxiety utilizing group systematic desensitization and relaxation training [dissertation]. Muncie, Ind: Ball State University, 1971.
- 215 Counts DK, Hollandsworth JG, Alcorn JD. Use of electromyographic biofeedback and cue-controlled relaxation in the treatment of test anxiety. *J Consult Clin Psychol* 1978; 46: 990-996.
- 216 Barrios BA, Ginter EJ, Scalise JJ, Miller FG. Treatment of test anxiety by applied relaxation and cue-controlled relaxation. *Psychol Rep* 1980; 46: 1287-1296.
- 217 Russell RK, Lent RW. Cue-controlled relaxation and systematic desensitization versus nonspecific factors in treating test anxiety. *J Counseling Psychol* 1982; 29: 100-103.
- 218 Sherman AR, Mulac A, McCann MJ. Synergistic effect of self-relaxation and rehearsal feedback in the treatment of subjective and behavioral dimensions of speech anxiety. *J Consult Clin Psychol* 1974; 42: 819-827.
- 219 Snyder AL, Deffenbacher JL. Comparison of relaxation as self-control and systematic desensitization in the treatment of test anxiety. *J Consult Clin Psychol* 1977; 45: 1202-1203.
- 220 Deffenbacher JL, Mathis H, Michaels AC. Two self-control procedures in the reduction of targeted and nontargeted anxieties. *J Counseling Psychol* 1979; 26: 120-127.
- 221 Marchetti A, McGlynn FD, Patterson AS. Effects of cue-controlled relaxation, a placebo treatment, and no treatment on changes in self-reported and psychophysiological indices of test anxiety among college students. *Behav Modif* 1977; 1: 47-72.
- 222 Kirkland K, Hollandsworth JG. Effective test taking: skills-acquisition versus anxiety-reduction techniques. *J Consult Clin Psychol* 1980; 48: 431-439.
- 223 Freeling NW, Shemberg KM. The alleviation of test anxiety by systematic desensitization. *Behav Res Ther* 1970; 8: 293-299.
- 224 Aponte JF, Aponte CE. Group preprogrammed systematic desensitization without the simultaneous presentation of aversive scenes with relaxation training. *Behav Res Ther* 1971; 9: 337-346.
- 225 Ricketts MS, Galloway RE. Effects of three different one-hour single session treatments for test anxiety. *Psychol Rep* 1984; 54: 115-120.
- 226 Johnson SM, Sechrest L. Comparison of desensitization and progressive muscle relaxation in treating test anxiety. *J Consult Clin Psychol* 1968; 32: 280-286.
- 227 Broota A, Sanghvi C. Efficacy of two relaxation techniques in examination anxiety. Special section: relaxation techniques and psychological management. *J Pers Clin Studies* 1994; 10: 29-35.
- 228 Comer JF. Meditation and progressive relaxation in the treatment of test anxiety [dissertation]. Lawrence: University of Kansas, 1977.
- 229 Bedell JR. Systematic desensitization, relaxation-training and suggestion in the treatment of test anxiety. *Behav Res Ther* 1975; 14: 309-311.
- 230 Mason TW. Treatment of test anxious students by applied cue-controlled relaxation, cue-controlled relaxation, applied relaxation, relaxation placebo and no-treatment [dissertation]. Lubbock: Texas Tech University, 1982.
- 231 Hurwitz LR, Kahane J, Mathieson C. The effect of EMG biofeedback and progressive muscle relaxation on the reduction of test anxiety. *Educ Psychol Res* 1986; 6: 291-298.
- 232 Sahasi G, Mohan D, Kacker C. Effectiveness of yogic techniques in the management of anxiety. *J Personality Clin Stud* 1989; 5: 51-55.
- 233 Castaneda R, Sussman N, Levy R, et al. A review of the effects of moderate alcohol intake on psychiatric and sleep disorders. *Recent Dev Alcohol* 1998; 14: 197-226.
- 234 Kushner MG, Abrams K, Borchardt C. The relationship between anxiety disorders and alcohol use disorders: a review of major perspectives and findings. *Clin Psychol Rev* 2000; 20: 149-171.
- 235 Kushner MG, Mackenzie TB, Fiszdon J, et al. The effects of alcohol consumption on laboratory-induced panic and state anxiety. *Arch Gen Psychiatry* 1996; 53: 264-270.
- 236 Abrams K, Kushner MG, Medina KL, Voight A. The pharmacologic and expectancy effects of alcohol on social anxiety in individuals with social phobia. *Drug Alcohol Depend* 2001; 64: 219-231.
- 237 MacDonald AB, Stewart SH, Hutson R, et al. The roles of alcohol and alcohol expectancy in the dampening of responses to hyperventilation among high anxiety sensitive young adults. *Addict Behav* 2001; 26: 841-867.
- 238 Lehman CL. Effects of alcohol consumption on laboratory-induced panic in patients with panic disorder: the role of alcohol outcome expectancy [dissertation]. Boston, Mass: Boston University, 2000.
- 239 Rimm D, Briddell D, Zimmerman M, Caddy G. The effects of alcohol and the expectancy of alcohol on snake fear. *Addict Behav* 1981; 6: 47-51.
- 240 Himle JA, Abelson JL, Haghhighatgou H, et al. Effect of alcohol on social phobic anxiety. *Am J Psychiatry* 1999; 156: 1237-1243.
- 241 Naftolowitz DF, Vaughn BV, Ranc J, Tancer ME. Response to alcohol in social phobia. *Anxiety* 1994; 1: 96-99.
- 242 Thyer BA, Curtis GC. The effects of ethanol intoxication on phobic anxiety. *Behav Res Ther* 1984; 22: 599-610.
- 243 Keane TM, Lisman SA. Alcohol and social anxiety in males: behavioral, cognitive, and physiological effects. *J Abnorm Psychol* 1980; 89: 213-223.
- 244 Thevos AK, Johnston AL, Latham PK, et al. Symptoms of anxiety in inpatient alcoholics with and without DSM-III-R anxiety diagnoses. *Alcohol Clin Exp Res* 1991; 15: 102-105.
- 245 Wetterling T, Junghans K. Psychopathology of alcoholics during withdrawal and early abstinence. *Eur Psychiatry* 2000; 15: 483-488.
- 246 Liappas J, Paparrigopoulos T, Tzavellas E, Christodoulou G. Impact of alcohol detoxification on anxiety and depressive symptoms. *Drug Alcohol Depend* 2002; 68: 215-220.
- 247 Driessen M, Meier S, Hill A, et al. The course of anxiety, depression and drinking behaviours after completed detoxification in alcoholics with and without comorbid anxiety and depressive disorders. *Alcohol* 2001; 36: 249-255.
- 248 Brown SA, Irwin M, Schuckit MA. Changes in anxiety among abstinent male alcoholics. *J Stud Alcohol* 1991; 52: 55-61.
- 249 Roberts MC, Emsley RA, Pienaar WP, Stein DJ. Anxiety disorders among abstinent alcohol-dependent patients. *Psychiatr Serv* 1999; 50: 1359-1361.
- 250 Uhde TW. Caffeine-induced anxiety: an ideal chemical model of panic disorder? In: Asnis GM, van Praag HM, editors. Panic disorder: clinical, biological, and treatment aspects. Detroit: Einstein Psychiatry, 1995: 181-205.
- 251 Bruce MS, Lader M. Caffeine abstinence in the management of anxiety disorders. *Psychol Med* 1989; 19: 211-214.
- 252 Smith GA. Caffeine reduction as an adjunct to anxiety management. *Br J Clin Psychol* 1988; 27: 265-266.
- 253 Salzer MS, Berenbaum H. Somatic sensations, anxiety, and control in panic disorder. *J Behav Ther Exp Psychiatry* 1994; 25: 75-80.
- 254 Uhde TW. Caffeine provocation of panic: a focus on biological mechanisms. In: Ballenger JC, editor. Neurobiological aspects of panic disorder. New York: Alan Liss, 1990: 219-242.
- 255 Uhde TW, Boulenger JP, Jimerson DC, Post RM. Caffeine: relationship to human anxiety, plasma MHPG and cortisol. *Psychopharmacol Bull* 1984; 20: 426-430.
- 256 Boulenger JP, Uhde TW, Wolff EA, Post RM. Increased sensitivity to caffeine in patients with panic disorders: preliminary evidence. *Arch Gen Psychiatry* 1984; 41: 1067-1071.
- 257 Bruce M, Scott N, Shine P, Lader M. Anxiogenic effects of caffeine in patients with anxiety disorders. *Arch Gen Psychiatry* 1992; 49: 867-869.
- 258 Newman F, Stein MB, Trettau JR, et al. Quantitative electroencephalographic effects of caffeine in panic disorder. *Psychiatry Res* 1992; 45: 105-113.
- 259 Bond AJ, Shine P, Bruce M. Validation of visual analogue scales in anxiety. *Int J Methods Psychiatr Res* 1995; 5: 1-9.
- 260 Totten GL, France CR. Physiological and subjective anxiety responses to caffeine and stress in nonclinical panic. *J Anxiety Disord* 1995; 9: 473-488.

- 261 Beck GJ, Berisford AM. The effects of caffeine on panic patients: response components of anxiety. *Behav Ther* 1992; 23: 405-422.
- 262 Tancer ME, Stein MB, Uhde TW. Lactic acid response to caffeine in panic disorder: comparison with social phobics and normal controls. *Anxiety* 1994-95; 1: 138-140.
- 263 Davis DD, Dunlop SR, Shea P, et al. Biological stress responses in high and low trait anxious students. *Biol Psychiatry* 1985; 20: 843-851.
- 264 Lee MA, Cameron OG, Greden JF. Anxiety and caffeine consumption in people with anxiety disorders. *Psychiatry Res* 1985; 15: 211-217.
- 265 Markus CR, Panhuysen G, Jonkman LM, Bachman M. Carbohydrate intake improves cognitive performance of stress-prone individuals under controllable laboratory stress. *Br J Nutr* 1999; 82: 457-467.
- 266 Wurtman RJ. Nutrients affecting brain composition and behavior. *Integr Psychiatry* 1987; 5: 226-257.
- 267 Parrott AC. Nesbitt's Paradox resolved? Stress and arousal modulation in cigarette smoking. *Addiction* 1998; 93: 27-39.
- 268 Jarvik ME, Henningfield JE. Pharmacological treatment of tobacco dependence. *Pharmacol Biochem Behav* 1988; 30: 279-294.
- 269 Parrott AC. Stress modulation over the day in cigarette smokers. *Addiction* 1995; 90: 233-244.
- 270 West R, Hajek P. What happens to anxiety levels on giving up smoking? *Am J Psychiatry* 1997; 154: 1589-1592.
- 271 Parker G, Hadzi-Pavlovic D. Is any female preponderance in depression secondary to a primary female preponderance in anxiety disorders? *Acta Psychiatr Scand* 2001; 103: 252-256.

(Received 1 Dec 2003, accepted 27 Feb 2004)

□