

# Preparing for out-of-season influenza epidemics when international travel resumes

Sheena G Sullivan

Influenza vaccination for travellers and the resources mobilised during the COVID-19 pandemic could mitigate the effects of an influenza resurgence



In this issue of the *MJA*, Marsh and colleagues<sup>1</sup> report that inter-seasonal influenza activity in New South Wales was unusually high during the 2018–19 summer, and provide compelling evidence for the role of international travel in seeding local influenza epidemics. During December 2018 – March 2019, people with notified influenza infections were 3.7 times as likely to report recent overseas travel as other people;

however, the odds were almost seven times as high for people with infections identified during the first two months of summer, suggesting that some of these cases initiated local outbreaks, with subsequent community transmission.<sup>1</sup>

This interpretation is consistent with previous findings that influenza circulation in Australia undergoes serial, local eliminations at the end of winter, with epidemics re-ignited by new viruses imported from overseas via international travel.<sup>2</sup> This phenomenon has never been better exemplified than during the coronavirus disease 2019 (COVID-19) pandemic. In early 2020, influenza activity was higher than normal for the inter-seasonal period, but plummeted with the introduction of COVID-19 restrictions in March 2020.<sup>3</sup> Introduction of new viruses was prevented by the closure of the Australian international border and the requirement that all returning travellers undergo 14 days of hotel quarantine. As a result, we have not confirmed the presence of influenza virus in any community samples sent to our reference laboratory since April 2020. It would seem that the virus has been locally eliminated, but the threat of its re-introduction looms with the imminent re-opening of our border.<sup>4</sup>

Throughout the pandemic, influenza A(H1N1)pdm09, A(H3N2), and B/Victoria have continued to circulate globally, with evidence of small, sporadic outbreaks, chiefly in tropical and subtropical regions of Asia and Africa (Box), where influenza seasonality is less marked than in temperate regions and year-round, low level activity is typical.<sup>5</sup> Marsh and colleagues<sup>1</sup> found that returned travellers with notified influenza had most frequently arrived from South-East Asia (42 of 168, 25%). However, current circulation patterns suggest that this region is not the most likely source of influenza importation into Australia. Our laboratory has confirmed the presence of influenza viruses in samples collected from travellers in hotel quarantine who have returned from South Asia, particularly those in the Howard Springs facility in the Northern Territory. As the influenza incubation period, infectious period, and serial interval are short,<sup>6</sup> the current duration of hotel quarantine (14 days) has prevented people leaving quarantine while still infectious, even in the context of chains of



transmission within travelling groups. However, if quarantine periods are reduced or eliminated, infected travellers may enter the community and instigate local outbreaks.

The report by Marsh and colleagues<sup>1</sup> indicates that, as borders open in late 2021, we should expect the introduction of influenza viruses into Australia and possibly intense out-of-season activity. Re-opening will precede our seasonal influenza vaccination campaign, which usually begins in April. To restrict the entry of influenza viruses, all returning travellers should be required to be vaccinated against influenza prior to travel, as recommended by the New South Wales Ministry of Health in response to the study by Marsh and colleagues.<sup>1</sup> Given the similarities in disease presentation, diagnosis, and management, the wealth of resources made available during the COVID-19 pandemic should also be leveraged to mitigate the consequences of influenza outbreaks. To avoid unclear messaging and to harmonise the response to all respiratory virus outbreaks, this would include testing for influenza as well as the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), especially in people in hotel quarantine, and sharing the resources of the special COVID-19 public health divisions that have been established in many jurisdictions.

Any resurgence of influenza will be most harshly felt by older people,<sup>7</sup> particularly those in aged care facilities, among whom influenza mortality is high<sup>8</sup> and the COVID-19 pandemic has already exerted a heavy toll.<sup>9</sup> It behoves us as a nation to protect our older citizens, and it reflects on us poorly when we fail to do so. In addition, influenza is likely to infect a considerable proportion of young children. In contrast to those infected with SARS-CoV-2, many children infected with influenza, especially those less than six months old, may need hospital care.<sup>7</sup> A resurgence of respiratory syncytial virus outbreaks in late 2020/early 2021 has already challenged some paediatric hospitals in Australia.<sup>10</sup> Increased severity of disease among patients co-infected with SARS-CoV-2 and influenza was reported in China in 2020,<sup>11</sup> and this could also have consequences for our healthcare system.

